

**Fig. 8.** Particulate  $^{234}\text{Th}$  ( $A_p$ , normalised to the dissolved activity  $A_d$ ) as a function of POC in suspended material collected near the Polar Front at 20-60m depth (open circles) and below 80m (closed circles), an example of the kind of data used to determine the POC/ $^{234}\text{Th}$  ratio on particles suspended at the depth where the export is to be determined. Summer expedition, ANT XIII/2 (Rutgers van der Loeff et al. 2002)


tained from a high-resolution transect of total  $^{234}\text{Th}$  in surface water alone as shown by the examples given above (Figs. 6,7). Another example is given by a transect from the SE Pacific (Fig. 9), where a strong export could be demonstrated near the ice edge, probably related to iron inputs (de Baar et al. 1999).

We conclude that the under-way analysis of total  $^{234}\text{Th}$  in surface waters, collected at high resolution from the ship's seawater supply, is a powerful tool to describe the geographical distribution of export production. Such a monitoring, supported by occasional measurements of activity-depth profiles and/or of the mixed layer depth, and linked with a satellite-based (SeaWiFS) monitoring of the distribution of chlorophyll (calibrated with shipboard observations, e.g. Bathmann et al. 1997), would provide a tool to estimate export production on a basin-wide scale.


### *The Fate of Exported Particles: Shallow Mineralization in the Weddell Sea*

In an area of the Weddell Sea with extremely low suspended particle load (61-67°S, Fig. 7) we found for the first time a situation with export from the surface mixed layer that was balanced by a release at only 150-350m depth (Fig. 10) implying a very shallow mineralization. This situation is extraordinary, as it is associated with a net uptake of  $\text{CO}_2$ , which is in part exported to greater depths through the formation of AABW, whereas no record of this export is retained in the underlying sediments (Usbeck et al. 2002; Hoppema et al. 1997).

This shallow mineralization is also reflected in the distribution of dissolved nutrients (Whitworth and Nowlin 1987), as shown in detail by inverse modelling of available hydrographic and nutrient data sets from the literature (Usbeck 1999; Usbeck et al. 2002; Schlitzer 2000, 2002). It is also found in the distribution of oxygen and  $\text{CO}_2$  (Hoppema et al. 1997), and in the distribution of another particle-reactive nuclide:  $^{210}\text{Pb}$  (Farley and Turekian 1990). The mineralization depth recorded with  $^{234}\text{Th}$  is somewhat shallower than expected from the distribution of these other tracers. This is probably related to the short half-life of  $^{234}\text{Th}$ , which implies that the  $^{234}\text{Th}$  release at 150-350m is asso-



**Fig. 9.** April 1995: Total  $^{234}\text{Th}/^{238}\text{U}$  ratios on a transect across the ACC in the SE Pacific (90°W, *Polarstern* Expedition ANT XII/4, redrawn from de Baar et al. 1999)



**Fig. 10.** Depth profiles of total  $^{234}\text{Th}/^{238}\text{U}$  ratios, showing remineralization maxima and negligible net export production below 350m in the central Weddell Sea (ANT XVI/3, redrawn from Usbeck et al. 2002)

ciated with the decomposition of the most labile fractions, whereas the distribution of the other tracers is determined by the cumulative decomposition of a wider range of fractions.

### $^{234}\text{Th}$ in the Nepheloid Layer

Near the seafloor a situation exists that is very similar to that in the surface ocean. In the benthic nepheloid layer (BNL),  $^{234}\text{Th}$  is adsorbed on resuspended particles. The resuspension-sedimentation cycle causes a depletion of total (dissolved + particulate)  $^{234}\text{Th}$  in a layer that is on the order of 100m thick. This depletion can be used to quantify the particle exchange between sediment and bottom water (Bacon and Rutgers van der Loeff 1989; Rutgers van der Loeff and Boudreau 1997). We have measured the distribution of  $^{234}\text{Th}$  in bottom waters on three transects across the Antarctic Circumpolar Current (ACC) in the southeast Atlantic (approx.  $0^\circ$  and  $40^\circ\text{E}$ , ANT XI/4) and in the southeast Pacific ( $90^\circ\text{W}$ , ANT XII/4) (Fig. 11). The distribution of particulate  $^{234}\text{Th}$  can be considered as a measure of particle load. Whereas in surface waters particulate  $^{234}\text{Th}$  is well correlated with al-

gal biomass and chlorophyll, in deep waters it depicts the intensity of the nepheloid layer. Relative to the low particle loads in the far south (central Weddell Sea), we observe enhanced particle loads associated with the major fronts in the ACC, probably as a result of high bottom water currents and of rugged topography.

In the entire area we observed only minimal depletion of  $^{234}\text{Th}$  with respect to  $^{238}\text{U}$  in the bottom water. This implies that the residence time of particles in the nepheloid layer is generally longer than a few weeks, the time scale that can be measured with this tracer. One station in the central Enderby basin with high suspended load in the bottom water formed a conspicuous exception with a depletion of 13% (Fig. 12). Apparently, a strong resuspension and partial sedimentation of the opal rich sediments had occurred here.


An essential parameter of the budget of  $^{234}\text{Th}$  in the BNL is the activity of  $^{234}\text{Th}$  on the resuspended material. The interaction between resuspension and bioturbation can be described in a model that can be calibrated with  $^{234}\text{Th}$  measurements. With this model (Rutgers van der Loeff and Boudreau 1997), turnover rates of other exchange processes at the sediment-water interface can be gauged with the  $^{234}\text{Th}$  tracer.

### Water Mass Tracers: $^{228}\text{Ra}$ , $^{227}\text{Ac}$

Radium and Actinium are relatively mobile elements with isotopes that are produced in the uranium decay series from highly insoluble parents. Consequently, they are very well suited to trace water masses after their contact with sediment surfaces. In the remote oceans like the Southern Ocean it is of particular importance to have a tool to trace the advection of water masses with a potential input of terrigenous micronutrients. In the framework of investigations on the sources of iron in the South Atlantic we have studied the distribution of the isotopes  $^{228}\text{Ra}$  and  $^{227}\text{Ac}$ .


#### $^{228}\text{Ra}$ : Tracer for Shelf Waters

$^{228}\text{Ra}$  is produced by decay of  $^{232}\text{Th}$ , which is ubiquitous in all sediments.  $^{228}\text{Ra}$  is therefore released by all sediments, both in the deep-sea and near-



**Fig. 11.** Transects of particulate  $^{234}\text{Th}$  on N-S sections across the ACC in the SE Pacific, the SE Atlantic and the SW Indian Ocean. South of the ACC, suspended loads in the bottom water are generally low.

## Enderby Basin, 60°S, 37°E



**Fig. 12.** An example of an exceptionally well-developed nepheloid layer in the center of the Enderby Basin. The 13% depletion of total  $^{234}\text{Th}$  ( $2.8 \text{ dpm cm}^{-2}$ ) implies an intensive exchange with surface sediments with an average residence time of about 12 days.

shore. The extended residence time of coastal water masses in shallow shelf seas allows the tracer to build up here far higher activities than are reached in near-bottom waters in the deep-sea.  $^{228}\text{Ra}$  is therefore especially well suited as a tracer of shelf waters. After the fundamental work of the GEOSECS expedition had established global distribution patterns of many radionuclide tracers including  $^{228}\text{Ra}$  (Li et al. 1980), the first transect across the ACC towards the Antarctic continent (Rutgers van der Loeff 1994) confirmed the expected enrichment of  $^{228}\text{Ra}$  on the Antarctic shelf. In the framework of the search for pathways of iron to the surface waters of the High Nutrient Low Chlorophyll (HNLC) Southern Ocean we have recently completed a detailed survey of  $^{228}\text{Ra}$  in the Atlantic sector of the Southern Ocean (Hanfland 2002). Gamma spectroscopy of radium fractions collected from surface waters on  $\text{MnO}_2$  adsorbents revealed the strong  $^{228}\text{Ra}$  sources both near South Africa (in the Agulhas Current, carrying signals obtained on the east-African coast), near the Antarctic continent (both near Neumayer station and

near the Antarctic Peninsula) and near the south American continent. In the Argentinean Basin, enhanced  $^{228}\text{Ra}$  activities were also found far offshore, indicating the far-reaching continental influence in the south Atlantic Gyre, in support of the studies of Li et al. (1980).

In the ACC proper, Hanfland (2002) mapped the distribution of  $^{228}\text{Ra}$  in the surface water with the far more sensitive  $^{228}\text{Th}$  ingrowth technique (after Li et al. 1980). The idea was, that if shelf-influenced water masses intrude into the frontal jets of the ACC, this ought to be visible in enhanced  $^{228}\text{Ra}$  levels. Radioactive decay of this 5.8-y half-life tracer cannot be significant during the transit across the Atlantic sector of the ACC, which can be estimated to be on the order of several months. But on several high-resolution sections across the ACC at  $0^\circ$  to  $20^\circ\text{E}$  Hanfland observed mostly very low  $^{228}\text{Ra}$  activities, discounting the role of this transport route for terrigenous material. As it can be expected that uptake and scavenging cause iron to be removed more rapidly than radium, Hanfland concluded that the shelf of South America, of the

Peninsula or of the south Sandwich Islands is not an important source for iron to alleviate the iron stress of plankton communities in the SE Atlantic (De Baar et al. 1995).

### *<sup>227</sup>Ac: Tracer for Deep Upwelling*

The application of this tracer is very similar to the previous tracer but with two major differences:

- the half-life of <sup>227</sup>Ac (21.8 y) is much longer (<sup>228</sup>Ra 5.8 y). The tracer is therefore suitable for transport processes on a timescale of decades rather than years.

- the activity of the parent nuclide of <sup>227</sup>Ac (<sup>231</sup>Pa) in sediments is dependent on water depth. As a consequence, deep-sea sediments are a far more important source for <sup>227</sup>Ac than shallow sediments.

Geibert (2001; Geibert et al. 2002) compared the release rate of <sup>228</sup>Ra and <sup>227</sup>Ac from marine sediments into the overlying water. This release rate depends not only on the activity of the mother nuclide in surface sediments, but also on the fraction of the daughter reaching the pore water, the adsorption equilibrium in the sediment pore water, and the bioturbation rate. Based on a model of Cochran and Krishnaswami (1980) for <sup>228</sup>Ra and an application of this model to <sup>227</sup>Ac by Nozaki et al. (1990) he shows that the <sup>227</sup>Ac/<sup>228</sup>Ra release rate ratio is indeed highly distinctive, changing from 0.026 for shelf sediments to 1.4 for the deep-sea. Consequently, <sup>227</sup>Ac can be considered as a specific tracer for contact with deep-sea sediments.

The first <sup>227</sup>Ac profiles from the Southern Ocean (Geibert 2001) show indeed this bottom source of <sup>227</sup>Ac, and can in principle be used to quantify mixing rates in the deep-sea. Similar studies have been performed based on the distribution of <sup>228</sup>Ra, but can now be extended to the longer time scale of <sup>227</sup>Ac. It is remarkable that these profiles show significant excess <sup>227</sup>Ac activities (i.e. in excess over the amount supported by <sup>231</sup>Pa in the water column) in surface waters south of the Polar Front. This implies that <sup>227</sup>Ac can be used here as a tracer for upwelling of water masses from the deep-sea. Upwelling rates in the Weddell Sea as based on these <sup>227</sup>Ac data (approx. 55 ma<sup>-1</sup>) are in line with earlier estimates based on heat budgets. The tracer appears to have a large potential for

studies of upwelling and diapycnal mixing rates in the ocean. The tracer may thus be used to monitor temporal changes in upwelling rates. If in the past two decades deep water production in the Weddell Sea had been smaller than usual, as has been hypothesised by Broecker et al. (1998), this would require a corresponding change in upwelling rate. If the circulation returned to normal, this might be detected as an increase in the <sup>227</sup>Ac levels in surface waters.

### Concluding Remarks

Natural radionuclides help us to study the transport of particles and water masses in the southern Atlantic Ocean.

Water masses that flow over continental shelf regions obtain a strong <sup>228</sup>Ra signal. Mostly very low <sup>228</sup>Ra activities in the frontal jets of the ACC show that such water masses cannot be responsible for the local fertilization in the central ACC with terrigenous trace substances like iron.

Upwelling of deep water south of the Polar Front can be followed with <sup>227</sup>Ac, a promising new tracer for water masses that have been in contact with deep-sea sediments.

Scavenging of particle-reactive isotopes helps us to calibrate particle fluxes. Export production, as measured with <sup>234</sup>Th, is absent in winter, shows large pulses in spring and continues with moderate rates during summer and autumn, with highest rates associated with the fronts of the ACC. South of the ACC, in the Weddell Gyre, we find a significant export production from the euphotic zone, but a highly efficient shallow remineralization in the depth zone of 150-350m, leaving only extremely low particle fluxes below 350m.

The low particle flux south of the ACC causes inefficient scavenging of <sup>230</sup>Th, observed in low <sup>230</sup>Th inventories and an accumulation of <sup>230</sup>Th in the water column. <sup>230</sup>Th is thus exported with AABW to the ACC where the surplus is removed by scavenging.

In the ACC, strong currents down to the sea-floor are responsible for resuspension of particles in the bottom waters, as observed by increases in turbidity and in particulate <sup>234</sup>Th. This causes a large-scale redistribution of sediments, with accu-

mulation rates that can exceed the local rain rates by an order of magnitude.  $^{230}\text{Th}$  is very suitable to correct for these focusing and winnowing effects, provided that the boundary scavenging of  $^{230}\text{Th}$  in the water column is taken into account.

Boundary scavenging is much more pronounced for  $^{231}\text{Pa}$  than for  $^{230}\text{Th}$ , making the Southern Ocean a sink for  $^{231}\text{Pa}$  produced further north in the Atlantic. The demonstration that this process continued during the last glacial (Yu et al. 1996) tells us that NADW transport continued during the LGM but not at what intensity. The lack of fractionation between Th and Pa in the opal-dominated Southern Ocean south of the Polar Front causes the  $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$  ratio here to have very limited value as proxy for (paleo)productivity.

### Acknowledgments

We are grateful for the assistance given by captains and crew of RV *Polarstern* on the many expeditions on which this paper is based. We wish to thank Dieter Fütterer for his full hearted support of the geochemistry group. M. Roy-Barman and R. Francois provided helpful reviews. This work was supported by the Deutsche Forschungsgemeinschaft (Sonderforschungsbereich 261). Data are available under [www.pangaea.de/Projects/SFB261](http://www.pangaea.de/Projects/SFB261).

### References

- Anderson HL, Francois R, Moran SB (1992) Experimental evidence for differential adsorption of Th and Pa on different solid phases in seawater. EOS 73 (43S, fall meeting), 270 p
- Anderson RF, Bacon MP, Brewer PG (1983a) Removal of  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  at ocean margins. Earth Planet Sci Lett 66: 73-90
- Anderson RF, Bacon MP, Brewer PG (1983b) Removal of  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  from the open ocean. Earth Planet Sci Lett 62: 7-23
- Andersson PS, Gustafsson O, Roos P, Broman D, Toneby A (2000) Particle mediated surface water export: comparison of estimates from  $^{238}\text{U}$ - $^{234}\text{Th}$  disequilibria and sediment traps in a continental shelf region. EOS AGU/ASLO Ocean Sciences Meeting, San Antonio, abstract only
- Asmus T, Frank M, Koschmieder C, Frank N, Gersonde R, Kuhn G, Mangini A (1999) Variations of biogenic particle flux in the southern Atlantic section of the Subantarctic Front during the late Quaternary: Evidence from sedimentary  $^{231}\text{Pa}_{\text{ex}}$  and  $^{230}\text{Th}_{\text{ex}}$ . Mar Geol 159: 63-78
- Bacon MP (1976) Applications of Pb-210/Ra-226 and Po-210/Pb-210 disequilibria in the study of marine geo-chemical processes. PhD thesis, Woods Hole Oceanographic Institution
- Bacon MP, Anderson RF (1982) Distribution of thorium isotopes between dissolved and particulate forms in the deep-sea. J Geophys Res 87: 2045-2056
- Bacon MP, Rosholt JN (1982) Accumulation rates of  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  and some transition metals on the Bermuda Rise. Geochim Cosmochim Acta 46: 651-666
- Bacon MP, Rutgers van der Loeff MM (1989) Removal of Thorium-234 by scavenging in the bottom nepheloid layer of the ocean. Earth Planet Sci Lett 92: 157-164
- Bacon MP, Spencer DW, Brewer PG (1976)  $^{210}\text{Pb}/^{226}\text{Ra}$  and  $^{210}\text{Po}/^{210}\text{Pb}$  disequilibria in seawater and suspended particulate matter. Earth Planet Sci Lett 32: 277-296
- Bacon MP, Huh C-A, Fleer AP, Deuser WG (1985) Seasonality in the flux of natural radionuclides and plutonium in the deep Sargasso Sea. Deep-Sea Res 32: 273-286
- Bathmann UV, Scharek R, Klaas C, Dubischar CD, Smetacek V (1997) Spring development of phytoplankton biomass and composition in major water masses of the Atlantic sector of the Southern Ocean. Deep-Sea Res II 44: 51-67
- Benitez-Nelson C, Buesseler KO, Rutgers van der Loeff MM, Andrews J, Ball L, Crossin G, Charette MA (2001) Testing a new small-volume technique for determining thorium-234 in seawater. J Radioanalytical Nuclear Chem 248: 795-799
- Broecker WS, Peng T-H (1982) Tracers in the Sea. Lamont- Doherty Geol Obs, Columbia University 690 p
- Broecker WS, Goddard J, Sarmiento JL (1976) The distribution of  $^{226}\text{Ra}$  in the Atlantic Ocean. Earth Planet Sci Lett 32: 220-235
- Broecker WS, Peacock SL, Walker S, Weiss R, Fahrback E, Schroeder M, Mikolajewicz U, Heinze C, Key R, Peng T-H, Rubin S (1998) How much deep water is formed in the Southern Ocean. J Geophys Res 103(C8): 15833-15843
- Buesseler K O (1998) The decoupling of production and particulate export in the surface ocean. Glob Biogeochem Cycl 12: 297-310
- Buesseler KO, Bacon MP, Cochran JK, Livingston HD

- (1992) Carbon and nitrogen export during the JGOFS North Atlantic Bloom Experiment estimated from  $^{234}\text{Th}$ : $^{238}\text{U}$  disequilibria. *Deep-Sea Res* 39: 1115-1137
- Buesseler KO, Andrews JA, Hartman MC, Belostock R, Chai F (1995) Regional estimates of the export flux of particulate organic carbon derived from thorium-234 during the JGOFS EqPac program. *Deep-Sea Res II* 42: 777-804
- Buesseler KO, Benitez-Nelson C, Rutgers van der Loeff MM, Andrews J, Ball L, Crossin G, Charette MA (2001) A comparison of methods with a new small-volume technique for thorium-234 in seawater. *Mar Chem* 74: 15-28
- Chase Z (2001) Trace elements as regulators (Fe) and recorders (U, Pa, Th, Be) of biological productivity in the ocean. PhD, Columbia University
- Chase Z, Anderson RF, Fleisher MQ, Kubik P (2002) The influence of particle composition on scavenging of Th, Pa and Be in the ocean. *Earth Planet Sci Lett* 204: 215-229
- Chase Z, Anderson RF, Fleisher MQ, Kubik P (2003) Scavenging of  $^{230}\text{Th}$ ,  $^{231}\text{Pa}$  and  $^{10}\text{Be}$  in the Southern Ocean (SW Pacific sector): The importance of particle flux and advection. *Deep-Sea Res II* 50: 739-768
- Chen JH, Edwards LR, Wasserburg GJ (1986)  $^{238}\text{U}$ ,  $^{234}\text{U}$  and  $^{232}\text{Th}$  in seawater. *Earth Planet Sci Lett* 80: 241-251
- Chung Y (1981)  $^{210}\text{Pb}$  and  $^{226}\text{Ra}$  distributions in the Circumpolar waters. *Earth Planet Sci Lett* 55: 205-216
- Chung Y, Applequist MD (1980)  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$  in the Weddell Sea. *Earth Planet Sci Lett* 49: 401-410
- Coale KH, Bruland KW (1985)  $^{234}\text{Th}$ : $^{238}\text{U}$  disequilibria within the California current. *Limnol Oceanogr* 30: 22-33
- Cochran JK (1992) The oceanic chemistry of the Uranium and Thorium-series nuclides. In: Ivanovich M and Harmon RS (eds) *Uranium-Series Disequilibrium: Applications to Earth, Marine, and Environmental Sciences*. 2nd edition, Clarendon Press, pp 334-395
- Cochran JK, Krishnaswami S (1980) Radium, Thorium, Uranium and  $^{210}\text{Pb}$  in deep-sea sediments and sediment pore waters from the North Equatorial Pacific. *Am J Sci* 280: 849-889
- Cochran JK, Buesseler KO, Bacon MP, Wang HW, Hirschberg DJ, Ball L, Andrews J, Crossin G, Fleer A (2000) Short-lived thorium isotopes ( $^{234}\text{Th}$ ,  $^{228}\text{Th}$ ) as indicators of POC export and particle cycling in the Ross Sea, Southern Ocean. *Deep-Sea Res II* 47: 3451-3490
- De Baar HJW, De Jong JTM, Bakker DCE, Löscher BM, Veth C, Bathmann U, Smetacek V (1995) Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean. *Nature* 373: 412-415
- De Baar HJW, de Jong JTM, Nolting RF, Timmermans KR, van Leeuwe MA, Bathmann U, Rutgers van der Loeff M, Sildam J (1999) Low dissolved Fe and the absence of diatom blooms in remote Pacific waters of the Southern Ocean. *Mar Chem* 36: 1-34
- Farley KA, Turekian KK (1990) Lead-210 in the circumpolar South Atlantic. *Deep-Sea Res* 37(12): 1849-1860
- Francois R, Bacon MP, Altabet MA, Labeyrie LD (1993) Glacial/Interglacial changes in sediment rain rate in the SW Indian sector of subantarctic waters as recorded by  $^{230}\text{Th}$ ,  $^{231}\text{Pa}$ , U, and  $\delta^{15}\text{N}$ . *Paleoceanography* 8: 611-629
- Francois R, Frank M, Rutgers van der Loeff MM, Bacon MP (2003)  $^{230}\text{Th}$ -normalization: An essential tool for interpreting sedimentary fluxes during the late Quaternary. *Paleoceanography*, submitted
- Frank M, Gersonde R, Mangini A (1999) Sediment redistribution,  $^{230}\text{Th}_{\text{ex}}$  - normalization and implications for the reconstruction of particle flux and export paleoproductivity. In: Fischer G, Wefer G (eds) *Use of Proxies in Paleoclimatology*. Springer, Berlin, pp 409-426
- Frank M, Gersonde R, Rutgers van der Loeff MM, Bohrmann G, Nürnberg CC, Kubik PW, Suter M, Mangini A (2000) Similar glacial and interglacial export bioproductivity in the Atlantic sector of the Southern Ocean: Multiproxy evidence and implications for glacial atmospheric  $\text{CO}_2$ . *Paleoceanography* 15: 642-658
- Friedrich J (1997) Polonium-210 und Blei-210 im Südpolarmeer: Natürliche Tracer für biologische und hydrographische Prozesse im Oberflächenwasser des Antarktischen Zirkumpolarstroms und des Weddellmeeres. *Ber Polarforsch* 235: 1-155
- Friedrich J, Rutgers van der Loeff MM (2002) A two-tracer ( $^{210}\text{Po}$ - $^{234}\text{Th}$ ) approach to distinguish organic carbon and biogenic silica export flux in the Antarctic Circumpolar Current. *Deep-Sea Res I* 49: 101-120
- Geibert W (2001) Actinium-227 as tracer for advection and mixing in the Deep-Sea. *Reports on Polar and Marine Research* 385, 112 p
- Geibert W, Usbeck R (2003) The adsorption of Thorium and Protactinium onto different particle types: Experimental findings. *Geochim Cosmochim Acta*, in press
- Geibert W, Rutgers van der Loeff MM, Hanfland C, Dauelsberg H-J (2002) Actinium-227 as a Deep-Sea Tracer: Sources, Distribution and Applications. *Earth Planet Sci Lett* 198: 147-165
- Gustafsson Ö, Düker A, Larsson J, Andersson P, Ingri

- J (2000) Functional separation of colloids and gravitoids in surface waters based on differential settling velocity: Coupled cross-flow filtration - split flow-thin cell fractionation (CFF-SPLITT). *Limnol Oceanogr* 45: 1731-1742
- Hanfland C (2002) Radium-226 and Radium-228 in the Atlantic sector of the Southern Ocean. Reports on Polar and Marine Research. PhD Thesis Univ Bremen, Germany 431, 135 p
- Henderson GM, Heinze C, Anderson RF, Winguth AME (1999) Global distribution of the  $^{230}\text{Th}$  flux to ocean sediments constrained by GCM modelling. *Deep-Sea Res I* 46: 1861-1893
- Hoppema M, Fährbach E, Schröder M (1997) On the total carbon dioxide and oxygen signature of the Circumpolar Deep Water in the Weddell Gyre. *Oceanol Acta* 20: 783-798
- Ku TL, Lin MC (1976)  $^{226}\text{Ra}$  distribution in the Antarctic Ocean. *Earth Planet Sci Lett* 32: 236-248
- Ku TL, Knauss KG, Mathieu GG (1977) Uranium in the open ocean: Concentration and isotopic composition. *Deep-Sea Res* 24: 1005-1017
- Kumar N, Gwiazda R, Anderson RF, Froelich PN (1993)  $^{231}\text{Pa}/^{230}\text{Th}$  ratios in sediments as a proxy for past changes in Southern Ocean productivity. *Nature* 362: 45-48
- Kumar N, Anderson RF, Mortlock RA, Froelich PN, Kubik P, Dittrich-Hannen B, Suter M (1995) Increased biological productivity and export production in the glacial Southern Ocean. *Nature* 378: 675-680
- Li Y-H, Feely HW, Toggweiler JR (1980)  $^{228}\text{Ra}$  and  $^{228}\text{Th}$  concentrations in GEOSECS Atlantic surface waters. *Deep-Sea Res* 27A: 545-555
- Löscher BM, de Jong JTM, de Baar HJW, Veth C, Dehairs F (1997) The distribution of Fe in the Antarctic Circumpolar Current. *Deep-Sea Res II* 44(1/2): 143-187
- Luo S, Ku T-L (1999) Oceanic  $^{231}\text{Pa}/^{230}\text{Th}$  ratio influence by particle composition and remineralization. *Earth Planet Sci Lett* 167: 183-195
- Marchal O, Francois R, Stocker TF, Joos F (2000) Ocean thermohaline circulation and sedimentary  $^{231}\text{Pa}/^{230}\text{Th}$  ratio. *Paleoceanography* 15: 625-641
- Nozaki Y, Horibe Y, Tsubota H (1981) The water column distributions of thorium isotopes in the western North Pacific. *Earth Planet Sci Lett* 54: 203-216
- Nozaki Y, Yamada M, Nikaido H (1990) The marine geochemistry of Actinium-227: Evidence for its migration through sediment pore water. *Geophys Res Lett* 17: 1933-1936
- Rutgers van der Loeff MM (1994)  $^{228}\text{Ra}$  and  $^{228}\text{Th}$  in the Weddell Sea. In: Johannessen OM, Muench RD, Overland JE (eds) *The Polar Oceans and their Role in Shaping the Global Environment: The Nansen Centennial Volume*. Geophysical Monograph 85, American Geophysical Union, pp 177-186
- Rutgers van der Loeff MM (2001) Uranium-Thorium decay series in the water column. In: Steele J, Thorpe S, Turekian K (eds) *Encyclopedia of Ocean Sciences*. Vol. MS 168, Academic Press
- Rutgers van der Loeff MM, Berger GW (1991) Scavenging and particle flux: Seasonal and regional variations in the South Ocean (Atlantic sector) *Mar Chem* 35: 553-567
- Rutgers van der Loeff MM, Berger GW (1993) Scavenging of  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  near the Antarctic Polar Front in the South Atlantic. *Deep-Sea Res I* 40: 339-357
- Rutgers van der Loeff MM, Boudreau BP (1997) The effect of resuspension on chemical exchanges at the sediment water interface - A modelling and natural radiotracer approach. *J Mar Syst* 11: 305-342
- Rutgers van der Loeff MM, Moore WS (1999) Determination of natural radioactive tracers. In: Grasshoff K, Kremling K, Ehrhardt M (eds) *Methods of Seawater Analysis*. Wiley-VCH, Weinheim, pp 365-397
- Rutgers van der Loeff MM, Westernströer U (2000) Export production measured through the  $^{234}\text{Th}/^{238}\text{U}$  disequilibrium in surface waters. In: Bathmann U, Smetacek V, Reinke M (eds) *The Expeditions ANTARKTIS XVI/3-4 of the RV Polarstern in 1999*. Rep Polar Res 364, pp 114-118
- Rutgers van der Loeff MM, Friedrich J, Bathmann UV (1997) Carbon export during the spring bloom at the southern Polar Front, determined with the natural tracer  $^{234}\text{Th}$ . *Deep-Sea Res II* 44: 457-478
- Rutgers van der Loeff MM, Buesseler KO, Bathmann U, Hense I, Andrews J (2002) Comparison of carbon and opal export rates between summer and spring bloom periods in the region of the Antarctic Polar Front, SE Atlantic. *Deep-Sea Res II* 49: 3849-3869
- Schlitzer R (2000) Applying the Adjoint Method for Global Biogeochemical Modeling. In: Kasibhatla P, Heimann M, Hartley D, Mahowald N, Prinn R, Rayner P (eds) *Inverse Methods in Biogeochemical Cycles*. AGU, pp 107-124
- Schlitzer R (2002) Carbon export fluxes in the Southern Ocean: Results from inverse modeling and comparison with satellite based estimates. *Deep-Sea Res II* 49: 1623-1644
- Scholten JC, Fietzke J, Vogler S, Rutgers van der Loeff MM, Mangini A, Koeve W, Stoffers P, Antia A, Neuer S, Waniek J (2001) Trapping efficiencies of



- sediment traps from the deep waster north Atlantic: The  $^{230}\text{Th}$  calibration. *Deep-Sea Res II JGOFS North Atlantic Synthesis* 48: 2383-2408
- Suman DO, Bacon MP (1989) Variations in Holocene sedimentation in the North American Basin determined from  $^{230}\text{Th}$  measurements. *Deep-Sea Res* 36: 869-878
- Usbeck R (1999) Modeling of marine biogeochemical cycles with an emphasis on vertical particle fluxes. *Ber Polarforsch* 332, 105 p
- Usbeck R, Rutgers van der Loeff MM, Hoppema M, Schlitzer R (2002) Shallow mineralization in the Weddell Gyre. *Geochem Geophys Geosyst* 3, 1: 10.1029/2001GC000182
- Usbeck R, Schlitzer R, Fischer G, Wefer G (2003) Particle fluxes in the ocean: Comparison of sediment trap data with results from inverse modeling. *J Mar Syst* 39: 167-183
- van Franeker J (1994) Sea-ice cover and icebergs. In: Bathmann UV, Smetacek V, de Baar HJW (eds) *The Expedition Antarktis X/6-8 of the research vessel Polarstern in 1992/1993*. *Ber Polarforsch* 135, AWI, Bremerhaven, pp 17-22
- Vogler S, Scholten J, Rutgers van der Loeff M, Mangini A (1998)  $^{230}\text{Th}$  in the eastern North Atlantic: The importance of water mass ventilation in the balance of  $^{230}\text{Th}$ . *Earth Planet Sci Lett* 156: 61-74
- Walter HJ, Rutgers van der Loeff MM, Höltzen H (1997) Enhanced scavenging of  $^{231}\text{Pa}$  relative to  $^{230}\text{Th}$  in the South Atlantic south of the Polar Front: Implications for the use of the  $^{231}\text{Pa}/^{230}\text{Th}$  ratio as a paleo-productivity proxy. *Earth Planet Sci Lett* 149: 85-100
- Walter HJ, Rutgers van der Loeff MM, Francois R (1999) Reliability of the  $^{231}\text{Pa}/^{230}\text{Th}$  activity ratio as a tracer for bioproductivity of the ocean. In: Fischer G, Wefer G (eds) *Use of Proxies in Paleooceanography - Examples from the South Atlantic*. Springer, Berlin, pp 393-408
- Walter HJ, Rutgers van der Loeff MM, Höltzen H, Bathmann U (2000) Reduced scavenging of  $^{230}\text{Th}$  in the Weddell Sea: Implications for paleoceanographic reconstructions in the South Atlantic. *Deep-Sea Res I* 47: 1369-1387
- Whitworth T, Nowlin WD (1987) Water masses and currents of the Southern Ocean at the Greenwich meridian. *J Geophys Res* 92(C6): 6462-6476
- Yu E-F, Francois R, Bacon MP (1996) Similar rates of modern and last-glacial ocean thermohaline circulation inferred from radiochemical data. *Nature* 379: 689-694
- Yu E-F, Francois R, Bacon MP, Honjo S, Flier AP, Manganini SJ, Rutgers van der Loeff MM, Ittekkot V (2001) Trapping efficiency of bottom-tethered sediment traps estimated from the intercepted fluxes of  $^{230}\text{Th}$  and  $^{231}\text{Pa}$ . *Deep-Sea Res I* 48: 865-889