

Chapter 2

Extrinsic and Intrinsic Factors Modulating Proliferation and Self-renewal of Multipotential CNS Progenitors and Adult Neural Stem Cells of the Subventricular Zone

SARA GIL-PEROTIN^{1,2} AND PATRIZIA CASACCIA-BONNEFIL¹

Introduction

Regulation of cell number in germinal zones of the nervous system is dependent on the interaction of extracellular signals with the “intrinsic” properties of the germinal cells that may vary depending on the developing stage of the organism. During early embryonic development, proliferation of cells occurs along the lumen of the developing neural tube, in an area defined as “the ventricular zone”. At this stage, cells proliferate very fast and characteristically give rise to identical daughter cells, via a process identified as “symmetric cell division” that allows for expansion of the primordial structures (Fig. 2.1A). As the organism develops, the need for “rapid expansion” decreases and new structures begin to form while still allowing for growth of the organism. Therefore, by mid-gestation, a second germinal zone arises, the subventricular zone (SVZ) and cells in this area acquire a modality of cell division characterized by the generation of two different daughter cells (“asymmetric cell division”): One with the ability to self-renew and the other one with the ability to differentiate into a specific lineage (Temple, 2001). In adult SVZ, the maintenance of homeostasis induces stem cells and multipotential progenitors to divide asymmetrically, unless a need for rapid expansion (e.g. repair after injury) induces the cells to shift to a symmetric modality of division (Fig. 2.1B).

Changes in the levels of the extracellular signals, alterations of their receptors or modification of the intracellular signaling molecules regulating proliferation during embryonic development, may result in abnormalities of

¹Department of Neuroscience and Cell Biology, UMDNJ R. Wood Johnson Medical School, 675 Hoes Lane, Piscataway, NJ 08854, USA

²Department of Comparative Neurobiology, Instituto Cavanilles de Biodiversidad y Biología evolutiva, 46980 Paterna, Valencia, Spain




FIGURE 2.1. Schematic representation of the distinct modalities of cell division. During development (A), the expansion of brain structures is first guaranteed by the rapid and symmetric non-terminal cell division. As new cell types are generated, the pattern of cell division becomes asymmetric. In the adult animal (B) it is likely that stem cells (B cells) in the remaining germinal zones such as the SVZ undergo asymmetric cell division to maintain a specific number of mother and daughter cells. “Multipotent progenitors (C cells) undergo a similar pattern of asymmetric cell division and generate A cells and oligodendrocyte progenitors (not shown), with the ability to divide following a symmetric division where both daughter cells exit from the cell cycle (Q) and differentiate. Note that upon injury, the need for expansion of the progenitor population leads to symmetric division and expansion of C cells that generate both neurons and oligodendrocytes.”

brain structures. Changes or modifications of extracellular or intracellular signals in adult neural stem cells, in contrast, may not affect the histoarchitecture of the brain, but affect the number of multipotential progenitors available for repair after injury. If proliferation is defective, a smaller number of cells will be available for repair, conversely, if proliferation proceeds uncontrolled, larger number of cells may result in hyperplastic foci and eventually lead to neoplastic transformation.

Since the responsiveness of neural stem cells and multipotential progenitors to extracellular signals is a dynamic process dependent on the developmental stage and on the regional localization of the cells, it becomes important to recognize that conclusions based on studies on embryonic stem cells may not be translated directly to adult neural stem cells. These temporally and developmentally restricted profiles of responsiveness to mitogenic and anti-mitogenic signals are determined by several parameters, including the bioavailability of extracellular signals, the presence of specific receptors, cross-talks among distinct signaling pathways and modulation of cell cycle regulatory molecules. All of these events can be affected or determined by specific genetic traits, expression of transcription factors and epigenetic modifications of chromatin components resulting in differences of gene expression that modulate the “context-specific” responsiveness of a stem cell.

It is worth mentioning that although the steady-state number of neural stem cells at any given stage of development is the result of the equilibrium between proliferation, differentiation, migration and survival of these cells, this chapter will focus only on the experimental evidence on extracellular factors and intracellular molecules affecting proliferation of neural stem cells and multipotent progenitors. This has been a very challenging task and although we have attempted to include several studies in this area, the overwhelming body of available literature has hindered our attempts to be exhaustive.

Extracellular Factors Affecting Proliferation

Basic Fibroblast Growth Factor (bFGF)

The Fibroblast Growth Factor (FGF) family includes a large number of ligands and receptors initiating signaling cascades that are critical for the early development of the organism (Burke *et al.*, 1998; Klint and Claesson-Welsh, 1999; Reuss and von Bohlen und Halbach, 2003). Of the different members of the FGF family of ligands, for instance, FGF8 is primarily involved in patterning of the midbrain and anterior forebrain (Mason *et al.*, 2000), FGF3 is important for the development of the ear (Represa *et al.*, 1991) and FGF2 is important for proliferation of neural stem cells and neurogenesis both in vitro (Reynolds and Weiss, 1992; Vescovi *et al.*, 1993; Vaccarino *et al.*, 1995; Kuhn *et al.*, 1997) and in vivo (Craig *et al.*, 1996; Tao *et al.*, 1996; Kuhn

et al., 1997; Wagner *et al.*, 1999). FGF2 (or bFGF) is expressed in the rodent brain at mid-gestation, from E11.5 to E17.5 in mice and from E13.5 to E19.5 in rats (Vacarino *et al.*, 1999b; Raballo *et al.*, 2000), during a period coincident with active neurogenesis (Bayer and Altman, 1991; Caviness *et al.*, 1995). Targeted deletion of *Fgf2* in mice results in a 50% reduction in the number of cortical neurons (Vacarino *et al.*, 1999b), thus suggesting a critical role for this ligand in neurogenesis.

The expression of the bFGF receptor, FGFR1 in the ventricular zone (Fig. 2.2), is observed at E8.5-9.5 (Orr-Urtreger *et al.*, 1991; Wanaka *et al.*, 1991) and it progressively decreases as neuroblasts begin to exit from the cell cycle and start differentiating (Raballo *et al.*, 2000). The phenotypic analysis of mutant mice with targeted deletions in the *Fgfr1* or *Fgfr2* receptors supports the idea that FGF signaling mediated by these two receptors, but not by FGFR3 and FGFR4, is critical for regulating proliferation and development of telencephalic structures (Yamaguchi *et al.*, 1994; Deng *et al.*, 1997; Partanen *et al.*, 1998; Xu *et al.*, 1998; Tropepe *et al.*, 1999).

The role of FGF receptor signaling in proliferation of neural progenitors and stem cells is also supported by a separate line of investigation on the effect of FGF2 administration at distinct developmental stages (Qian *et al.*, 1997; Kelly *et al.*, 2003). High doses of FGF2 intracerebrally injected during embryogenesis (E14 in mice), result in massive enlargement of the ventricles and aberrant proliferation and differentiation (Ohmiya *et al.*, 2001). However, low doses of recombinant FGF2 in rat embryos (Vacarino *et al.*, 1999a) or even in neonatal and adult rats (Tao *et al.*, 1996; Wagner *et al.*, 1999) enhance proliferation and neurogenesis.

Epidermal Growth Factor (EGF) Family

The epidermal growth factor (EGF) family of polypeptides includes EGF, transforming growth factor- α (TGF- α), heparin-binding EGF (HB-EGF) and related neuregulins. These polypeptides, produced by neurons and glial cells, play an important role in the development of the nervous system, by affecting proliferation, survival, migration and differentiation of neuronal and glial cells (Xian and Zhou, 1999). Neuregulins have been identified during the late embryonic development (Corfas *et al.*, 1995), and their receptors ErbB2 and ErbB4 are expressed in the E12 embryo (Kornblum *et al.*, 2000) as well as in embryonically derived neural stem cells (Calaora *et al.*, 2001). Although neuregulin signaling especially via the ErbB4 receptor is critical for migration of adult neuroblasts and possibly survival and neurogenesis (Calaora *et al.*, 2001; Anton *et al.*, 2004), the available experimental evidence does not support a role of this EGF family member in proliferation. TGF α , in contrast, is a potent mitogen and also the predominant form of EGF ligand expressed in the developing brain and in adult SVZ (Kornblum *et al.*, 1997). The importance of this ligand

in regulating proliferation of adult neural stem cells is supported by the phenotypic characterization of the TGF α null mice characterized by decreased number of mitosis exacerbated by senescence (Tropepe *et al.*, 1997).

Expression of EGFR *in vivo* occurs during late embryogenesis (Fig. 2.2) in the developing SVZ and follows the expression of FGFR1 in ventricular zone cells (Burrows *et al.*, 1997). Consistent with this temporal progression, cells isolated from embryonic mouse brain during early development (i.e. E10-E12) are FGFR1+, while those isolated at later developmental stages are both EGFR+ and FGFR1+ (Ciccolini and Svendsen 1998; Gritti *et al.*, 1999; Lillien and Raphael 2000). This expression pattern is also consistent with the distinct growth factor requirements of early embryonic stem cells for FGF2 and of the late embryonic stem cells for EGF and FGF2 (Tropepe *et al.*, 1999; Martens *et al.*, 2000).

The concept of temporal responsiveness to distinct growth factors determined by the sequential expression pattern of distinct receptor subunits is also supported by a comparative phenotypic analysis of the *Egfr* and *Fgfr* null mice. While the *Fgfr1* null mice are early embryonic lethal (Deng *et al.*, 1994; Yamaguchi *et al.*, 1994), the phenotype of the *Egfr* null mice is characterized by reduced cortical size at E18 (Threadgill *et al.*, 1995), and progressive neuronal degeneration during the postnatal period (Sibilia and Wagner, 1995; Sibilia *et al.*, 1998).

Despite the similar role as mitogens, FGF2 and EGF have been differentially implicated as modulators of lineage restriction and neurogenesis. It has been proposed that the differential role played by these factors depends on the segregation of the mitogenic effect on distinct cellular populations: EGF preferentially targeting the quiescent stem cells and FGF targeting the more committed neuroblasts (Morshead and van der Kooy, 1992; Morshead *et al.*, 1994). The co-expression of FGFR1 and EGFR within the same cell type at later stages of development, however, argues against this possibility (Gritti *et al.*, 1999). An alternative explanation for the differential effect exerted by these two growth factors is the possibility that through the activation of distinct tyrosine kinase receptors, they may affect distinct intracellular signaling effectors, or the kinetics of activation of specific signaling molecules (Lax *et al.*, 2002; Yamada *et al.*, 2004).

The idea that FGF2 and EGF may differentially affect the behavior of multipotent neural progenitors is also based on the evidence that two weeks of intraventricular administration of EGF to adult mice result in decreased neurogenesis and increased generation of astrocytes, while administration of FGF2 results in enhanced generation of neurons (Kuhn *et al.*, 1997). Similar data have been obtained by several other groups with the exception of a study, reporting similar effects of FGF2 or EGF treatment on adult neurogenesis (Craig *et al.*, 1996). The *in vivo* ultrastructural identification of adult SVZ cells affected by EGF intraventricular infusion, for instance, clearly

demonstrated the ability of EGF to affect the modality of cell division of the transit amplifying C cell population from asymmetric to symmetric, and to restrict neuroblast formation (Doetsch *et al.*, 2002a). Similarly, intrastriatal infusion of TGF alpha in an animal model of Parkinson's disease (Cooper and Isacson, 2004) induces the formation of clusters of GFAP⁻/nestin⁺ cells along the lateral wall of the ventricle, very likely representing an expansion of the transit amplifying C cells (Cooper and Isacson, 2004). Finally, in vitro studies in cultured neurospheres show increased astrocyte generation in response to EGF and enhanced neuronal differentiation in response to FGF2 (Whittemore *et al.*, 1999; Jori *et al.*, 2003). Together, these data identify EGF signaling as permissive for astrocytic but not neuronal differentiation and FGF signaling as permissive for the neurogenic fate in the adult SVZ.

Insulin-Like Growth Factor-1 (IGF1)

Insulin growth factor peptides 1 and 2 are members of a family of insulin-related peptides originally identified by their ability to stimulate growth of chondrocytes (Laron, 2001). IGF1 is secreted by many tissues and its function varies according to the site of secretion and the presence of its receptors. The expression of its receptors is highly conserved throughout evolution (Garofalo and Rosen, 1988). IGF1R, in particular, is expressed in the embryonic brain and co-localizes with the expression of FGFR and EGFR in cells of selective germinal zones (Bondy *et al.*, 1990; Garcia-Segura *et al.*, 1991; Kar *et al.*, 1993). The phenotype of mice with targeted deletion in the *Igf1* gene or in the *Igf1R* gene is severely hypomorphic, with a clear decrease in brain size (Baker *et al.*, 1993; Liu *et al.*, 1993; Beck *et al.*, 1995), thus suggesting IGF1R function as critical for brain development. Conversely, transgenic mice over-expressing IGF1 show a generalized increase in cell number and corresponding increase in brain size (Carson *et al.*, 1993). The role of IGF1 in neurogenesis is still controversial. While in vitro studies on embryonic and adult stem cells suggest a role in neuronal differentiation of EGF-responsive stem cells (Arsenijevic and Weiss 1998; Arsenijevic *et al.*, 2001), other reports underline its role as survival factor for FGF responsive stem cells (Drago *et al.*, 1991) and studies on freshly isolated PSA-NCAM⁺ cells describe IGF1 as both survival and mitogenic factor for EGF responsive cells (Gage *et al.*, 2003). Its function in oligodendrocytes and myelination is better characterized. In addition, it has been recently suggested that IGF1 also favors the commitment of adult neural stem cells towards the oligodendrocytic phenotype (Hsieh *et al.*, 2004). More studies on the in vivo function of IGF1 will be necessary to decipher the multiple roles played by this factor in the adult SVZ.

Neurotrophins

The neurotrophin family is composed of several trophic factors including the originally discovered founding member nerve growth factor (NGF) (reviewed in Aloe, 2004), and the related brain-derived neurotrophic factor (BDNF), neurotrophin-3, -4 and -5 (NT-3, NT-4, NT-5) (Leibrock *et al.*, 1989; Hohn *et al.*, 1990; Maisonpierre *et al.*, 1990; Berkemeier *et al.*, 1991; Hallbook *et al.*, 1991). Neurotrophins bind to two classes of receptors: tyrosine kinase receptors (Trk A, B and C) and a low affinity neurotrophin receptor called p75 that is structurally related to the TNFR superfamily (Barker, 2004). Each ligand binds with the highest affinity to a specific tyrosine kinase receptor (i.e. TrkA and NGF, TrkB and BDNF, TrkC and NT-3), while all the neurotrophins can bind to the low-affinity p75 (Chao, 2003). The complexity of the system is enhanced by the presence of alternatively spliced isoforms for TrkB and TrkC that may differ in case of the presence of specific catalytic domains (Huang and Reichardt, 2003; Teng and Hempstead, 2004).

In the adult SVZ p75 immunoreactivity is confined to proliferating cells. The majority of the p75⁺ cells are identified also by EGFR immunoreactivity. Few p75⁺ cells in the SVZ are also nestin⁺, but the majority of them does not colabel with GFAP or PSA-NCAM (Giuliani *et al.*, 2004), thus suggesting that p75 is mainly expressed in the fast-proliferating cell population. Interestingly, no TrkA receptor expression is detected in the periventricular region by *in situ* hybridization (Anderson *et al.*, 1995) or immunohistochemistry (Giuliani *et al.*, 2004; Fiore *et al.*, 2005), while the full-length and truncated form of TrkB receptors are both present. Truncated TrkB is confined to the ependymal cell layer and to choroid plexus, while full length TrkB expression is more widespread (Anderson *et al.*, 1995). Therefore, it is not surprising that BDNF is the primary neurotrophin-affecting neurogenesis in the adult SVZ (Kirschenbaum and Goldman, 1995). *In vivo* infusion of BDNF in the lateral ventricle of the rat adult brain enhances BrdU incorporation in the SVZ and is associated with an increased number of neurons migrating to the olfactory bulb (Zigova *et al.*, 1998). Since the BrdU⁺ cells are also p75⁺ and TrkB⁻ (Pencea *et al.*, 2001), it is thought that the proliferative effect of BDNF is mediated by signaling through the low affinity p75 receptor in the fast proliferating cell population. The effect of BDNF on adult neurogenesis (i.e. the generation of neurons from the multipotential stem cells), in contrast, is a TrkB-dependent event, and is likely directed on PSA-NCAM⁺ neuroblasts. In agreement with this model, treatment of embryonic and adult-derived neurospheres with neurotrophins affects differentiation, but not proliferation of TrkB⁺ cells (Ahmed *et al.*, 1995; Benoit *et al.*, 2001). BDNF also acts as permissive factor for the maturation and survival of neuroblasts generated from the SVZ (Kirschenbaum and Goldman 1995). A recent study on the effect of BDNF on GABAergic interneurons derived from the SVZ has shown that the sequential activation of p75 and then TrkB signaling pathways is critical for the

development of the dendritic arbor (Gascon *et al.*, 2005). Together, these studies support a p75-mediated effect in cell proliferation and a TrkB-mediated effect in neurogenesis of adult neural stem cells.

Ephrins

Ephrins are cell-surface-tethered ligands for Eph receptors, a family of tyrosine kinase receptors. The functional complex ephrin/Eph is involved in several processes, including the formation and guidance of growth cones from differentiating neurons and the induction and maturation of neuronal spines (Palmer and Klein, 2003). There are two subclasses of ephrin ligands: type A (ephrinA) are GPI-linked membrane proteins while type B (ephrinB) are transmembrane proteins (Orioli and Klein, 1997; O'Leary and Wilkinson, 1999). The ephrin family of receptors (Eph) can also be subdivided into class A (EphA) and B (EphB) on the basis of structural similarities in the extracellular domain and on their ability to preferentially bind to specific ligands. Type A ligands (ephrinA) bind to EphA receptors, and type B ligands (ephrinB) bind to EphB receptors, although the EphA4 receptor can bind to both types of ligands (Kullander and Klein, 2002) and ephrinA5 can also bind to the EphB2 receptor (Himanen *et al.*, 2004; Pasquale, 2004). During development, this signaling system modulates attraction/repulsion, cell adhesion and cell migration (Klein, 2004). In addition, a possible direct or indirect role for ephrinB1 in neurogenesis is suggested by its expression in neuroepithelial cells in the VZ at the onset of neocortical neurogenesis and its persistence throughout the neurogenetic period (Stuckmann *et al.*, 2001). It has been suggested that ephrinB1 plays a role in affecting the responsiveness of neuroepithelial cells to other cues and also to favor migration of newly generated neurons towards their targets.

In adult SVZ, both ephrin ligands (ephrin-B2,B3 and A5) and ephrin receptors (EphB1-B3, EphA4) are expressed in specific subpopulations of cells (Conover *et al.*, 2000). Intraventricular infusion of ephrin B2 or of the EphB2 ectodomain dramatically disrupts neuroblast migration and increases proliferation in the SVZ, thus resulting in the formation of regions of localized hyperplasia (Conover *et al.*, 2000).

Therefore, ephrins/Eph complexes act as environmental cues for migration processes, axonal pathfinding and topographic mapping during development, although they can also modulate proliferation and guided migration of neurons in the adult SVZ.

The Tgfbeta Family, Including Bone Morphogenetic Protein (BMP)

Transforming growth factor beta (TGFbeta) signaling controls several intracellular processes including proliferation, apoptosis, differentiation and

lineage specification. TGFbeta ligands bind to serine-threonine kinase receptors (type I and II) on the cell surface and the signal is mediated by a heterogeneous group of proteins called Smads (Shi and Massague, 2003).

The TGFbeta family of cytokines comprises two subfamilies, TGFbeta/Activin/Nodal subfamily and BMP (Bone Morphogenetic Protein)/GDF (Growth and Differentiation Factor)/MIS (Muellerian Inhibiting Substance) subfamily. TGFbeta cytokines are expressed in the CNS of the developing rodent (Flanders *et al.*, 1991; Millan *et al.*, 1991; Schmid *et al.*, 1991) in regions where neuronal differentiation occurs. In fact, TGFbeta2 *in vitro* induces cell cycle exit and differentiation of precursor cells (Mahanthappa and Schwarting, 1993; Constam *et al.*, 1994; Kane *et al.*, 1996). The BMP family includes a group of dorsal morphogens whose effect is pleiotropic and ranges from the induction of a dorsal fate in cells of the developing neural tube (Shah *et al.*, 1996; Liem *et al.*, 1997; Panchision *et al.*, 2001), to the suppression of differentiation and maintenance of self-renewal in embryonic stem cells (Ying *et al.*, 2003), from the down-regulation of EGFR expression in embryonic progenitors (Lillien and Raphael, 2000), to the induction of apoptosis (Graham *et al.*, 1996). In addition, BMP signaling has been implicated in neurogenesis (Liem *et al.*, 1995; Reissmann *et al.*, 1996; Li *et al.*, 1998; Panchision and McKay, 2002) as well as in gliogenesis (Gross *et al.*, 1996), and also to favor the commitment to the astrocytic lineage at the expenses of neurogenesis and oligodendroglialogenesis (Grinspan *et al.*, 2000; See *et al.*, 2004). The effect of BMPs on astroglialogenesis is dependent on cross-talks among distinct signaling pathways and involves the activation of critical signaling molecules, including SMADs and STATs (Nakashima *et al.*, 1999a). Since SMADs are downstream of BMP signaling and STATs are downstream of LIF signaling, it is the interaction between these two signaling pathways that appears to be critical for astroglialogenesis. Intriguingly, however, an alternative pathway of activation of STATs by BMP receptor signaling (Rajan *et al.*, 2003) has been suggested. According to this model, STAT activation is mediated by a serine-threonine kinase (called FRAP) that becomes activated upon binding of BMP4 to its receptor (Rajan *et al.*, 2003).

Besides their role in development, BMPs favor astroglialogenesis also in the adult animal (Lim *et al.*, 2000; Panchision *et al.*, 2001). Indeed, BMP2 and 4 and cognate receptors are expressed in the adult SVZ where they favor the astrocytic phenotype of adult neural stem cells (Lim *et al.*, 2000) and possibly modulate the cell cycle length of migrating neuroblasts (Coskun and Luskin, 2001). Noggin, a BMP antagonist expressed by the ependymal cells, promotes neurogenesis by counteracting the effect of the BMPs on astroglialogenesis (Lim *et al.*, 2000).

Thus, the activation of the TGFbeta signaling modulates both the decision of a cell to exit from the cell cycle and the commitment to an astrocytic fate.

Ciliary Neurotrophic Factor (CNTF) and Leukemia Inhibitory Factor (LIF)

Ciliary neurotrophic factor (CNTF) and leukemia inhibitory factor (LIF) are two neuroregulatory cytokines which play a major role in the developing nervous system. They both exhibit broad structural similarities (Bazan, 1991) and share signaling components (Gearing *et al.*, 1991, 1992; Gearing and Bruce, 1992; Ip *et al.*, 1992, 1993) among each other and with other members of the family, including interleukin-6, oncostatin or cardiotrophin 1. CNTF is widely expressed within the nervous system (Ip *et al.*, 1993; Ip and Yancopoulos, 1996; Ip, 1998) and has been implicated in fate choice decision and survival of sensory, sympathetic, ciliary and motor neurons (Sleeman *et al.*, 2000; Turnley and Bartlett 2000).

LIF and CNTF share a common receptor, gp130 (Davis *et al.*, 1993; Ip and Yancopoulos, 1996; Nandurkar *et al.*, 1996). Specificity of the signaling response is achieved by selective binding of the ligand with specific receptor components. LIF signaling requires dimerization of the LIF receptor subunit beta (LIFRbeta) with gp130; while CNTF requires trimerization of its cognate receptor subunit (CNTFRalpha) with LIFRbeta and gp130 (Fig. 2.2). CNTFRalpha is expressed in embryonic neural precursor cells (Ip *et al.*, 1993; Lachyankar *et al.*, 1997) and in neurons and astrocytes of the adult central nervous system (Ip *et al.*, 1993; MacLennan *et al.*, 1996; Lee *et al.*, 1997a,b; Kirsch *et al.*, 1998; Dallner *et al.*, 2002), including the subventricular zone (Seniuk-Tatton *et al.*, 1995).

In vitro studies on embryonic stem cells suggest a role for CNTF/LIF signaling in maintaining pluripotency (Conover *et al.*, 1993) and preventing differentiation (Pennica *et al.*, 1995) or even promoting survival (De Felici and Dolci, 1991; Pesce *et al.*, 1993). The phenotype of the *CNTF* null mice, however is relatively normal and exhibits motor neuron losses only later in life, thus arguing against a major role played by this cytokine during development (Masu *et al.*, 1993). The phenotypes of the *CNTFR* $-/-$ (DeChiara *et al.*, 1995), of the *LIFR* $-/-$ (Li *et al.*, 1995) or the *gp130* $-/-$ (Nakashima *et al.*, 1999a) mice, in contrast, are characterized by a profound motor neuron defect at birth, thus supporting the notion that CNTF is critical for survival and viability of motor neurons (Sendtner *et al.*, 1994; Ip 1998). The neuroprotective effect of CNTF is also observed in vivo, as demonstrated by the intracerebral administration of this cytokine in animal models of Huntington's disease (Anderson *et al.*, 1996; Emerich *et al.*, 1996) and in injured dopaminergic neurons after transection of the nigrostriatal pathway (Hagg and Varon, 1993).

The detection of CNTF receptors during embryonic development and in adult neural germinal zones raises the possibility that CNTF/LIF family members play a role in regulating proliferation and fate choice of neural stem cells. Treatment of neural embryonic progenitors and stem cells with these cytokines suggests a critical role in astroglial differentiation (Bonni *et al.*, 1997;

Rajan and McKay 1998; Park *et al.*, 1999; Galli *et al.*, 2000; Morrow *et al.*, 2001). Signaling through STAT3, a transcription factor downstream of the LIF/gp130 receptor signaling pathway is critical for the expression of GFAP (Bonni *et al.*, 1997) a fact also supported by the severely perturbed astroglialogenesis in *LIFR* $-/-$ mice (Koblar *et al.*, 1998). The role of STAT in astrocytic differentiation has been the subject of several studies. It has been shown that GFAP promoter activation requires the assembly of a complex including SMADs, STATs, the co-activator CBP and the histone acetyl-transferase p300 (Nakashima *et al.*, 1999b). Over-expression of neurogenin, a neuronal-specific basic HLH factor, disrupts this “gliogenic” complex by sequestration of the CBP/p300 component from STATs and thus prevents GFAP promoter activation (Sun *et al.*, 2001). The presence of histone acetyl transferase p300 in the transcriptional complex suggests that activation of the astrocytic program of differentiation necessitates changes in chromatin conformation. Besides acetylation of nucleosomal histones, the GFAP promoter is also regulated by a switch in the methylation of specific lysine residues on nucleosomal histones (Song and Ghosh, 2004). The “switch” from a silencing methylation on lysine 9 to an activating methylation on lysine 4 of histone H3 is affected by the presence of FGF2 and results in an open chromatin conformation in the promoter region, thus facilitating binding of transcriptional activators such as STATs and SMADs (Song and Ghosh 2004). CNTF signaling has also been implicated in oligodendrocytic maturation (Barres *et al.*, 1996; Marmur *et al.*, 1998) and neuronal differentiation (Ernsberger *et al.*, 1989; Saadat *et al.*, 1989; Ip *et al.*, 1994; Rudge *et al.*, 1996; Ezzeddine *et al.*, 1997; Lachyankar *et al.*, 1997). In the adult forebrain, signaling through the CNTFR/LIFR/gp130 complex is responsible for the maintenance of EGF-responsive neural stem cells (Fig. 2.3). CNTF treatment of SVZ-derived cells *in vitro*, increases self-renewal and expansion (Shimazaki *et al.*, 2001) and *in vivo*, it enhances proliferation of the EGF-responsive population (Shimazaki *et al.*, 2001; Chojnacki *et al.*, 2003). The effect of CNTF/LIF signaling on proliferation and self-renewal can be explained in terms of receptor cross-talks. Proliferation could be consequent to the effect of CNTF on Notch1 signaling (Chojnacki *et al.*, 2003), while self-renewal could be due to the effect of LIF on differentiation inhibitors, such as the Ids, downstream of BMP receptor signaling (Ying *et al.*, 2003).

Notch1

Notch is a cell-surface receptor activated by contact with a member of the DSL family of ligands (Delta, Serrate, Lag2). Upon ligand activation, the Notch receptor is cleaved and its intracellular domain (Notch ICD) is released into the cytosol, translocates into the nucleus where it activates the transcription of CSL/CBF and induces the expression of HES genes that have been described as basic HLH transcription factors with the ability to

inhibit neuronal differentiation (Lindsell *et al.*, 1996; Weinmaster, 2000). Therefore, Notch signaling during development has been linked to inhibition of differentiation (Artavanis-Tsakonas *et al.*, 1995). Although the role of Notch in timing of cell fate specification and differentiation is supported by several studies (Yun *et al.*, 2002), the persistence of Notch1 and Jagged 1 expression in the adult SVZ (Stump *et al.* 2002) suggests that these molecules may also modulate the behavior of pluripotential progenitors and adult stem cells. The depletion of neural stem cells in the *Notch1*^{-/-} mice (Hitoshi *et al.*, 2002) indicates a role for Notch in promoting self-renewal at the expenses of neurogenesis. However, transient Notch activation induced by administration of Notch ligand results in severe decrease of the neurogenic potential paralleled by increased gliogenesis (Morrison *et al.*, 2000b). These apparently contradictory results can be reconciled by evoking the importance of spatial and temporal cues on the responsiveness of progenitor cells to Notch signaling. Indeed, expression of active Notch at midgestation inhibits proliferation and decreases the generation of neurons (Chambers *et al.*, 2001). At later stages, however, Notch ICD promotes proliferation and gliogenesis (Gaiano *et al.*, 2000; Chambers *et al.*, 2001). Thus, similar to what was described for BMP and CNTF, the same signal can result in maintenance of stem-like cells or gliogenesis, depending on the cellular context.

Sonic Hedgehog

Sonic hedgehog (Shh), is a very well characterized morphogen expressed at high levels in cells of the ventral telencephalon at embryonic day 11.5 (E11.5) and maintained throughout development (Dahmane and Ruiz-i-Altaba 1999; Wallace 1999; Wechsler-Reya and Scott 1999). Shh has been implicated in several aspects of CNS development such as proliferation (Marti *et al.*, 1995; Roelink *et al.*, 1995; Chiang *et al.*, 1996; Ericson *et al.*, 1996) and cell fate determination (Zhu *et al.*, 1999). It has also been shown to exert opposing actions to BMP2 in embryonic cortical progenitors (Machold *et al.*, 2003, Viti *et al.*, 2003b). Mice, bearing conditional null alleles of both *Shh* and its receptor *Smoothed*, have a dramatic reduction in the number of neural progenitors in the SVZ, possibly resulting from reduced proliferation and increased apoptosis (Machold *et al.*, 2003).

Recent studies on the adult SVZ in postnatal and adult mice have identified the Shh responsive SVZ cells as the GFAP⁺ B cells and the EGF responsive transit amplifying progenitors C cells (Palma *et al.*, 2005). The *in vitro* data in SVZ cultures treated with Shh do not support a direct effect of this molecule on proliferation, although they do suggest a synergistic effect with EGF (Palma *et al.*, 2005). Similarly, the increased number of neurospheres formed by embryonic stem cells pre-treated with Shh and cultured in the presence of EGF has been ascribed to the up-regulation of EGFR level (Viti *et al.*, 2003b). The lack of proliferation or differentiation in the adult SVZ after intrastriatal injection of a myristoylated form of Shh

(Charytoniuk *et al.*, 2002) is consistent with the *in vitro* data. However, the decreased proliferation observed in the SVZ after administration of the Shh antagonist cyclopamine suggests a more complex role for this molecule (Palma *et al.*, 2005). Although the role of Shh in survival of SVZ cells has not been addressed, it is likely to play a role in modulating the responsiveness of neural stem cells to other signaling molecules regulating cell number (Fig. 2.3).

Wnt

The behavior of cells in the developing nervous system is tightly regulated by the highly conserved family of Wnt signaling molecules. Wnt proteins can either be secreted or located at the cell surface and may interact with a family of cell surface receptors in the Frizzled family (Ikeya *et al.*, 1997; Yoshikawa *et al.*, 1997; Hall *et al.*, 2000). Binding of the ligand to the




FIGURE 2.2. Extracellular Receptors in embryonic and adult neural stem cells. Schematic representation of the major subtypes of extracellular receptors observed during embryonic development (panel A) and in the adult SVZ (panel B). Note that during early embryonic development (upper cell in panel A), only FGF and LIF receptors are expressed, but at later stages cells become also responsive to BMPs, Shh and EGF (lower cell in panel A). In the adult SVZ (panel B), a differential pattern of receptor expression is observed. The relatively quiescent B cells are responsive to BMPs, Shh and neurotrophins ephrins, while the transit amplifying progenitors express the receptors for EGF, FGF, CNTF, and Notch (B).

receptor transduces a signal which involves inactivation of the GSK-3 kinase and the accumulation of the transcriptional regulator beta catenin. Of the several Wnt family members, analysis of the phenotype of mice with targeted deletions in specific genes has revealed the critical importance of Wnt1, Wnt3a and Wnt7a in the developing nervous system (Megason and McMahon, 2002). Cell proliferation is commonly regulated by Wnt signaling and expansion of the CNS fails in *Wnt1* mutants (reviewed in Logan and Nusse, 2004) Over-expression of Wnt7a in embryonic stem cells increases proliferation and self-renewal both in vivo and in vitro and further promotes maturation of cortical progenitors by inducing the expression of EGFR (Viti *et al.*, 2003b).

Recently, the role of Wnt and its downstream-signaling molecule beta catenin has been explored in neural stem cells. Transgenic mice over-expressing beta catenin have grossly enlarged brains that could not be simply explained in terms of mitogenic effect or decreased apoptosis (Chenn and Walsh, 2002). Rather, it appears that beta catenin affects the decision of progenitors to exit from the cell cycle and this, in turn, results in loss of growth control (Chenn and Walsh, 2002; Zechner *et al.*, 2003). However, in other cellular systems such as embryonic stem cells, beta catenin favors neurogenesis (Otero *et al.*, 2004). This cell context role of β catenin has been linked to the presence of FGF2 (Israsena *et al.*, 2004). In the presence of FGF2, beta catenin contributes to the maintenance of a proliferative state (Viti *et al.*, 2003), while in the absence of FGF2, it enhances neuronal differentiation by forming transcriptionally active complexes on neurogenic promoters (Israsena *et al.*, 2004; Otero *et al.*, 2004; Logan and Nusse, 2004). A better understanding of the role of Wnt pathway in neural stem cell biology will be a very important and critical step for the design of stem cell-based therapies.

Hypoxia-Induced Growth Factors

Ischemia and cerebral injury stimulate neurogenesis in neuroproliferative regions of the adult brain, including SVZ and the hippocampal DG (Gould and Tanapat, 1997; Parent *et al.*, 1997; Liu *et al.*, 1998; Takagi *et al.*, 1999; Gu *et al.*, 2000; Magavi *et al.*, 2000; Jin *et al.*, 2001; Yoshimura *et al.*, 2001; Zhang *et al.*, 2001). Concomitantly to ischemic injuries, expression of some factors increases (Kawahara *et al.*, 1999; Marti, 2004):

(a) Erythropoietin EPO

Erythropoietin (EPO) is a pleiotropic-inducible molecule produced by the kidney and whose function was first described as the regulator of red blood cell production (Carnot and Deflandre, 1906) by promoting erythrocyte survival in the bone marrow (Koury and Bondurant, 1990a; 1990b; Yousoufian *et al.*, 1993; Fisher, 2003). EPO is also a key example of a gene that is regulated in an oxygen-dependent manner and, thus, its expression is

induced when the oxygen levels are reduced (Wenger, 2002). Recently, EPO and its receptor EPOR have also been detected in the developing CNS, thus suggesting a possible role in neural development (Buemi *et al.*, 2002; Liu *et al.*, 1994; Juul *et al.*, 1998,1999). Indeed, mice with an *Epor* targeted deletion (*Epor*^{-/-}) (Lin *et al.*, 1996), are characterized by the severe reduction in the number of neural progenitor cells and increased apoptosis (Yu *et al.*, 2002).

The observation that embryonic precursors in the CNS proliferate and differentiate more in response to lowered oxygen (Morrison *et al.*, 2000a; Studer *et al.*, 2000) suggests that perhaps they could play a similar role in adult neural stem cells. In vitro studies on cultured neural stem cells are consistent with the idea that increased EPO gene expression results in increased adult neurogenesis (Shingo *et al.*, 2001). Furthermore, intraventricular infusion of EPO in mice favors the migration of newly generated neurons to the olfactory bulb and the effect is blocked by anti-EPO antibodies (Shingo *et al.*, 2001). Together these data suggest that EPO can negatively regulate proliferation of stem cells while favoring the differentiation towards the neuronal lineage.

(b) Vascular Endothelial Growth Factor VEGF

The vascular endothelial growth factor (VEGF) is a hypoxia-inducible secreted protein (Wenger, 2002) that regulates endothelial cell growth and differentiation and is also a survival factor for endothelial cells (Risau, 1997). The loss of a single allele in the mouse results in death during embryogenesis, due to vascular defects (Ferrara *et al.*, 1996). In the nervous system, VEGF is expressed during development (Breier *et al.*, 1992), and is related to the EPO-induced response to hypoxic insults in the brain as a target for the hypoxia inducible transcription factor (HIF-1) (Marti, 2004). VEGF has neurotrophic and neuroprotective effects on distinct types of neurons (Silverman *et al.*, 1999; Sondell *et al.*, 1999, 2000; Jin *et al.*, 2000a, 2000b; ; Matsuzaki *et al.*, 2001) and its receptor VEGFR2/flk-1 is expressed in neural progenitor cells (Yang and Cepko, 1996; Jin *et al.*, 2002b), thus suggesting a possible role in neurogenesis. In the adult murine brain, administration of exogenous VEGF increases proliferation (Fig. 2.3) of neuronal precursors in the SVZ by modulating cell division rather than survival (Jin *et al.*, 2002b). Finally, in cultures from the neonatal SVZ, treatment with FGF2 increases the expression of VEGFR2/flk-1, and in turn, treatment with VEGF enhances the chemotactic response of FGF2-stimulated progenitors, thus suggesting a synergistic effect of these two factors on migration (Zhang *et al.*, 2003).

(c) Heparin-Binding HB-EGF

Heparin-binding EGF-like growth factor (HB-EGF) is a mitogenic and chemotactic glycoprotein that contains an EGF-like domain and acts

through several receptors, including ErbB1, ErbB4, and heparin sulfate proteoglycans. Although the targeted deletion of *HB-EGF* in mice affects mainly heart and skin development (Yamazaki *et al.*, 2003), the expression in neurons and glial cells throughout the brain suggests a role in the CNS (Mishima *et al.*, 1996; Hayase *et al.*, 1998; Nakagawa *et al.*, 1998). As for EPO and VEGF, the expression of HB-EGF in the brain is increased by ischemia and results in neuroprotection (Kawahara *et al.*, 1999). In addition, HB-EGF enhances neurogenesis *in vitro*, in neonatal cerebellar cultures (Opanashuk and Hauser 1998) and embryonic mouse neurons exposed to hypoxic conditions (Jin *et al.*, 2002a). *In vivo*, intraventricular infusion HB-EGF enhances neurogenesis in the adult SVZ (Jin *et al.*, 2002a) and restores neurogenesis to young adult levels when administered to aged mice in combination with FGF2 (Jin *et al.*, 2003).

Extracellular Matrix and Cell–Cell Contact

It has been proposed that the maintenance of neural stem cells in the adult brain is favored by the presence of extracellular conditions creating a “niche” that favors the preservation of an undifferentiated and proliferative state (Doetsch, 2003; Alvarez-Buyilla and Lim, 2004). The concept of a “niche” including components of the extracellular matrix, is quite attractive and has also been described in the hematopoietic system (Mercier *et al.*, 2002). Remarkably, several components identified in the extracellular matrix of the SVZ (Gates *et al.*, 1995) have been proven effective in modulating the responsiveness to mitogens (i.e. FGF2, EGF) or to morphogens (i.e. Shh, Wnt, BMPs). For instance, ECM molecules such as Tenascin C and chondroitin sulfate proteoglycans, present in the late embryonic SVZ and persist in the adult brain (Garcion *et al.*, 2004), modulate the sensitivity to other extracellular signals at several developmental stages. This effect could be due to indirect binding to other matrix components or to direct interaction with specific cell surface receptors. In the tenascin null mice the responsiveness of embryonic stem cells to FGF2 is dramatically reduced, while the sensitivity to BMP4 is increased (Garcion *et al.*, 2004). Given the previously discussed antagonistic role of BMP and FGF2 on EGFR expression (Lillien and Raphael, 2000), it is not surprising that tenascin loss of function results in decreased proliferation of SVZ cells and delayed EGFR expression.

Another component of the ECM, the glycosaminoglycan heparin sulfate, has also been shown to promote the action of FGF2 in embryonically derived cells (Chipperfield *et al.*, 2003), although it inhibits the response to this same factor in cells derived from the adult brain (Leventhal *et al.*, 1999; Shen *et al.*, 2004). These data corroborate and support the idea that the extracellular matrix is a critical component of the niche and that it may affect stem cell behavior by modulating the responsiveness to other extracellular cues and possibly affecting intracellular signals.

Another essential component of the neural stem cell niche is the vascular compartment. In the developing CNS, the embryonic neural stem cells in the VZ have been shown to produce vascular endothelial growth factor (VEGF) which is known to contribute to the neovascularization of the area. A more direct evidence that endothelial cells enhance proliferation and neurogenesis of embryonic and adult neural stem cells is provided by co-culture experiments (Leventhal *et al.*, 1999). Explants of adult SVZ cultured in the presence of endothelial cells express higher levels of the neurotrophin BDNF (Shen *et al.*, 2004). Time-lapse video recording of dividing clones of neural stem cells, grown in the presence of endothelial cells, indicates that co-culture conditions tend to favor the symmetric modality of cell division (Shen *et al.*, 2004). Therefore, proliferation of the neural stem cells seems to be affected by a wide range of molecular signals, including the production of soluble factors (i.e. VEGF, BDNF), the cross talk with the wnt/beta catenin signaling pathway and/or with the Notch signaling pathway (Temple, 2001, Shen *et al.*, 2004).

Neurotransmitters

(a) Dopamine

Dopamine is a neurotransmitter produced by neurons in the substantia nigra, ventral tegmental area and preoptic area. It is involved in numerous brain processes and contributes to integration of cortical information underlying motor, limbic and cognitive aspects of behavior (Nieoullon, 2002).

Besides its function as neuromodulator, dopamine also plays a role in neurogenesis during development. The D1 and D2-receptors are expressed in the striatal VZ and have been shown to play opposing roles in favoring (D2) or inhibiting (D1) cell cycle progression in the lateral ganglionic eminence (Jung and Bennett, 1996). The effect of D1-receptor activation is dominant over the effect of the D2 receptor and results in an overall reduction of cells entering S-phase (Ohtani *et al.*, 2003). The role of D3 receptor signaling is not well established, although it is expressed in the proliferative neuroepithelium and persists postnatally in the subventricular zone (Diaz *et al.*, 1997). Administration, either in vivo or in vitro of D3-receptor agonists, increases the proliferative rate of neural stem cells and the number of cells expressing neuronal markers (Pilon *et al.*, 1994; Coronas *et al.*, 2004; Van Kampen *et al.*, 2004). This effect is mediated by MAPK activation, a pathway also activated by BDNF to affect neurogenesis (Zigova *et al.*, 1998; Pencea *et al.*, 2001). Given the dual relationship between dopamine receptor activation and BDNF expression (Guillin *et al.*, 2001, 2003; Kuppens and Beyer, 2001; Sokoloff *et al.*, 2002;), it is likely that they synergize in promoting neurogenesis.

(b) Serotonin (5-HT)

Serotonin (5-HT) is produced by neurons of the raphe nucleus in the brain stem and modulates sensorimotor control, cognition and mood (Struder and

Weicker, 2001b, 2001a). In addition to modulating synaptic function in the adult brain, 5-HT also controls important functions in brain development such as neurite outgrowth, cell survival and synaptogenesis (Gaspar *et al.*, 2003).

The role of serotonin in neurogenesis is suggested by studies on the class of antidepressants called “Serotonin Selective Re-uptake Inhibitors (SSRI)”. Stress is known to inhibit neurogenesis by elevating the levels of gluco-corticoids (Moghaddam *et al.*, 1994; Stein-Behrens *et al.*, 1994). SSRI anti-depressants reverse the effect of stress and increase proliferation and differentiation of newly formed cells into neurons in the hippocampus (Malberg *et al.*, 2000; Santarelli *et al.*, 2003). Since serotonin receptors (5-HT1A and HT2C) are expressed in the SVZ, it is not surprising that systemic administration of various agonists increases proliferation of cells in this brain region (Banar *et al.*, 2004). Intriguingly, like for dopamine, the effects of serotonin on neurogenesis seem to be related to BDNF signaling, thus suggesting that the effect of the distinct classes of neurotransmitters is possibly linked to the presence of neurotrophins (Mattson *et al.*, 2004).

(c) Opioids

Opioid peptides are known to act as neurotransmitters or neuromodulators in the adult nervous system. They act through three cognate receptors: μ , δ , κ (Dhawan *et al.*, 1996) that are also expressed in the SVZ (Zagon and McLaughlin, 1986; Stiene-Martin *et al.*, 2001). Blockade of opioid receptors enhances cell proliferation, while their activation induces an anti-proliferative effect (Hauser *et al.*, 1996). Although this effect was originally attributed to a fourth opioid receptor ζ (Zagon *et al.*, 1991), it is likely that the opioid effect on neurogenesis is a μ -mediated effect since the μ receptor is widely expressed postnatally in neuroproliferative regions (Stiene-Martin *et al.*, 2001).

Hormones

(a) Thyroid Hormone

T3 constitutes the active ligand of the thyroid hormone (TH). The expression of TH receptors in the brain varies according to the cell type, region and age as it clearly shows a spatial-temporal patterning during development (Bradley *et al.*, 1992) and adulthood (Puymirat *et al.*, 1991). Besides the well-established role of TH in maturation of oligodendrocytes (Baas *et al.*, 1997; Baumann and Pham-Dinh, 2001), the presence of its receptors in the adult brain also led to investigate a possible effect in neurogenesis. Indeed, hypothyroid rats showed increased proliferation in the SVZ and olfactory bulb, while hyperthyroid rats showed reduced proliferation and increased tendency to differentiate into oligodendrocytes (Fernandez *et al.*, 2004). Co-administration of thyroid hormone with retinoic acid results in a net increase of proliferation in SVZ and enhanced neurogenesis (Giardino *et al.*, 2000).

(b) Estrogens

The role of sex steroids in neurogenesis has been suggested by the existence of a gender bias in hippocampal-dependent tasks (Roof *et al.*, 1993; Frye *et al.*, 2000; Conrad *et al.*, 2003). At the cellular level, these differences are correlated with the proliferative effects of estrogens in the hippocampus (Tanapat *et al.*, 1999, 2005). Estrogens can bind to two types of receptors, called alpha and beta. Both receptors have been detected in several brain regions throughout development (Shughrue *et al.*, 1990). Both receptors are also present in the ventricular wall of the embryonic neural tube as well as in the adult brain (Brannvall *et al.*, 2002), but the functional role of estrogens at distinct stages of development is quite distinct. While estrogen treatment potentiates the mitogenic effect of EGF in embryonic neural stem cells, it antagonizes the EGF effect in adult neural stem cells, by upregulating the cell cycle inhibitor p21Cip/Waf1 (Brannvall *et al.*, 2002).

(c) Prolactin

Prolactin is a hormone that increases during pregnancy and at postpartum, signaling lactation. Prolactin stimulates the production of neuronal progenitors in the SVZ (Bridges and Grattan, 2003; Shingo *et al.*, 2003). The increased neurogenesis results in the formation of new neurons in the olfactory bulb (Shingo *et al.*, 2003), and is possibly related to the enhanced olfactory capability of the mother.

Others

(a) Amyloid Precursor Protein and Amyloid Peptide

The amyloid precursor protein (APP) is a type I transmembrane protein with unknown physiological functions. Its soluble-secreted form (sAPP), present in normal brain tissue (Palmer *et al.*, 1989), has biological activities resembling a growth factor and increases the *in vitro* proliferation of embryonic neural stem cells (Ohsawa *et al.*, 1999). The soluble sAPP binds to EGFR⁺ cells in the adult SVZ and *in vitro*, EGF induces the secretion of soluble APP (sAPP) by SVZ-derived cells. Intriguingly, sAPP infusions into the lateral ventricle enhances proliferation of the EGF-responsive progenitors and increases the cell number (Caille *et al.*, 2004). In pathological conditions such as Alzheimer's disease, however, neurons are exposed to the amyloid beta-peptide (A β), a self-aggregating neurotoxic protein. This peptide, in contrast to sAPP, has been shown to impair neurogenesis in the SVZ of adult mice and in human cortical neural precursor cells (Fig. 2.3). Amyloid beta peptide treatment suppresses both proliferation and differentiation of neural progenitors and induces apoptosis, associated with a disruption of calcium regulation. The cumulative result of these effects is a severe depletion of neurons possibly contributing to the olfactory and cognitive deficits observed in Alzheimer's disease (Haughey *et al.*, 2002).




FIGURE 2.3. **Schematic view of a sagittal section of the adult brain.** In red are some of the extracellular signals that inhibit proliferation and favor the exit from the cell cycle. In green are the extracellular signals that promote proliferation and increase neurogenesis.

Intracellular Signals Affecting Proliferation

Although it is often assumed that experimental results obtained in stem cells isolated in the developing animal can be extrapolated to the behavior of stem cells in the mature CNS, a large number of studies support the concept of intrinsic differences in distinct neural stem cell populations, depending on their location and birthdate (Temple 2001). The existence of temporally regulated changes intrinsic to the cell is suggested by studies of in vitro time-lapse videos of isolated stem cells. These studies have shown that cells maintained in the same culture conditions can first give rise to neurons and then to glia (Qian *et al.*, 1998, 2000).

These “intrinsic differences” may result from genetic differences and epigenetic modifications affecting the pattern of gene expression in a given cell population. Consistent with this interpretation, genetic profiling of embryonic and adult hematopoietic stem cells has identified a relatively small subset of commonly expressed genes and an even smaller number of genes shared with neural stem cells (Ivanova *et al.*, 2002). Changes in gene expression may also result from differences in the extrinsic signaling pathways whose cross talk affects the length of the cell cycle (T_c) and/or the probability of progenitor cells to re-enter the cell cycle or become quiescent (Nowakowski *et al.*, 2002). In this respect, it has been shown that cells in the embryonicVZ undergo a progressive increase in the length of T_c and that an increased proportion of these cells leaves the cell cycle with each cell division (Takahashi *et al.*, 1996). Both these events are likely to be modulated by the expression levels of cell cycle regulatory molecules and other transcription

factors (Tarui *et al.*, 2005). Thus, progressive changes in the expression of cell cycle genes modify the cycle kinetics and the relative proportion of proliferating cells within each population, depending on the developmental stage and cellular context.

Studies on the cell cycle kinetics of progenitors/neural stem cells during embryonic development, for instance, have reported increased cell cycle length with increasing embryonic age (Fig. 2.4) and a switch of cell division from symmetric and rapid ($T_c = \sim 17.6$ hr) at E11, to asymmetric and slower ($T_c = \sim 26.5$) at E14 (Tropepe *et al.*, 1999). Differences in cell division persist in the adult forebrain subependyma (Fig. 2.4), and at least two distinct populations of proliferating cells have been identified (Morshead *et al.*, 1998). One population, the constitutively proliferating population, has a T_c of 12.7 hr (Morshead and van der Kooy, 1992) and corresponds to the transit amplifying progenitor population (Doetsch *et al.*, 2002a), also called the “C cell type” (Doetsch *et al.*, 1997). The other population has a much longer cell cycle duration ($T_c \sim 15$ d or more) and corresponds to the quiescent cell population (Morshead *et al.*, 1994, 1998), of adult “stem cells” also called “B cell type” (Garcia-Verdugo *et al.*, 1998).

Lengthening of the cell cycle time is thought to be a function of an increase in the duration of G1 as the rest of the cell cycle parameters remain relatively




FIGURE 2.4. Lengthening of the cell cycle duration in stem cells during development. Note that during the early stages of development (A), the cell cycle duration (T_c) is very fast, possibly allowing for expansion. Around E11 the T_c is 17 hours and the modality of cell division primarily symmetric. As the organism develops and neurogenesis begins (E14) the cell cycle time increases to 26 hours and the modality of division becomes asymmetric. In the adult SVZ (B) two main cell types have been identified. The relatively quiescent B cells has a very long T_c (15 days) and has been proposed to be the precursor of the rapidly expanding population of C cells, characterized by a short cell cycle time (12 hrs) and the ability to give rise to neuroblasts and oligodendrocyte progenitors that become quiescent (Q).

constant over time (von Waechter and Jaensch, 1972; Caviness *et al.*, 1995). Therefore, it is likely that the expression of cell cycle regulatory molecules in distinct cell populations accounts for differences in cell cycle kinetics. In agreement with this model, studies on cell cycle length in the neonatal rat brain (Schultze and Korr, 1981; Menezes *et al.*, 1995; , 1998; Smith and Luskin, 1998) have indicated that differences in cell cycle kinetics between cells in the neonatal anterior SVZ (that have a fast cell cycle time) and the migratory cells in the RMS with a slower kinetics of cell division (Smith and Luskin, 1998), correlate with the levels of expression of the G1 inhibitor p19INK4d (Coskun and Luskin, 2001).

Indeed cell cycle length and the probability to exit from the cell cycle are both affected by cell cycle regulatory molecules and transcription factors whose expression can be modulated by genetic factors, epigenetic modifications of chromatin and by the integration of extracellular signals.

Cell Cycle Regulatory Molecules

In order to discuss intracellular mechanisms of proliferation of CNS progenitors and neural stem cells, it becomes critical to introduce the molecules regulating the progression from G1 into the S phase of the cell cycle. Progression through G1 is regulated by the ordered synthesis, assembly and activation of distinct cyclin-CDK enzymatic complexes (Dyson, 1998; Nevins, 2001). Two main enzymatic activities have been described: CDK4, acting in early-mid G1; and CDK2, acting in late G1, very close to the entry into the S replicative phase (Sherr, 1994; Sherr and Roberts, 1999). These two activities differ in terms of substrate specificity and modality of regulation. CDK4, for instance, is positively regulated by cyclin D and is inhibited by members of the INK4 (INhibitors of CDK4) family. CDK2, in contrast, is positively regulated by cyclin E and negatively regulated by the Kips (Kinase Inhibitory Proteins). The main substrates of cyclinD/CDK4/6 complexes are proteins of the Rb family (including pRb, p107 and p130). INK4 proteins prevent their phosphorylation, thus allowing them to sequester E2F and blocking the transcription of E2F-responsive genes that are responsible for driving the cell into S-phase (Kastan *et al.*, 1992). Besides the role of Rb as growth-inhibitory pathway, another important cell cycle checkpoint acting at the G1 phase is mediated by the p53 tumor-suppressor gene (Paggi *et al.*, 1996; Mundle and Saberwal, 2003). We shall now review literature pertinent to the expression patterns of these cell cycle regulatory molecules in the central nervous system, with a special emphasis on their possible functional role in the SVZ.

(a) Rb Family

The Retinoblastoma gene family is composed of three members of closely related proteins characterized by a “pocket” domain pRb, p107

and p130/Rb2 (Brehm *et al.*, 1998; Luo *et al.*, 1998; Magnaghi-Jaulin *et al.*, 1998). These molecules play a critical role in eukaryotic cell cycle progression as negative regulators of proliferation. The retinoblastoma gene product pRb, in its hypophosphorylated state, binds to members of the E2F family of transcription factors, converting them to active transcriptional repressors, by recruiting histone deacetylases (Dyson, 1998). Phosphorylated pRb in contrast, is unable to bind to E2F, the repression is relieved and results in the transcription of genes involved in DNA-replication (Fig.2.5) and nucleotide biosynthesis (Beijersbergen *et al.*, 1994; Ginsberg *et al.*, 1994). Distinct members of the Rb family show association with specific members of the E2F family and pRb preferentially binds to E2F-1, -2 and -3 while p107 and p130 preferentially bind to E2F-4 and -5 (Lees *et al.*, 1992; Li *et al.*, 1993). In addition, p107 and p130 can also bind to cyclin/CDK2 complexes (Gill *et al.*, 1998; Callaghan *et al.*, 1999; Ferguson and Slack, 2001).

The expression profile of the “pocket proteins” in the brain has a characteristic cellular and temporal pattern. While pRb is found in both dividing precursor cells and postmitotic neurons during embryogenesis, p107 expression is restricted to the ventricular zone and is rapidly down-regulated at the onset of differentiation (Jiang *et al.*, 1997; Yoshikawa 2000). P130 is expressed mainly in post-mitotic differentiated cells (Clarke *et al.*, 1992; Jacks *et al.*, 1992; Lee *et al.*, 1994). Consistent with the temporal pattern of expression, targeted deletions in the *Rb* locus result in embryonic lethality (Cobrinik *et al.*, 1996; Lee *et al.*, 1996), while mice with deletions in *p107* or *p130* develop normally (Vanderluit *et al.*, 2004). The expression pattern of




FIGURE 2.5. Molecular control of cell cycle entry. The G1/S transition of the cell cycle is regulated by the enzymatic activity of cyclin/CDK complexes. The resulting increased phosphorylation of the tumor suppressor gene pRb (or other members of the pocket protein family) induces the release of transcription factors of the family E2F/DP and allows the transcription of genes involved in S phase entry. The main inhibitors (INK4 and KIP family members) and activators (cyclin D and E) of the cyclin/CDK complexes active at the G1/S transition are shown.

p107 persists in the adult SVZ, where it is expressed in small clusters of cells around the ventricular wall (Vanderluit *et al.*, 2004). Mice lacking p107 exhibit increased proliferation of the fast proliferating population, but also increased self-renewal of neural stem cells, as indicated by the ability of cells derived from the SVZ of p107 null mice, to generate a larger number of secondary neurospheres than wild type mice (Vanderluit *et al.*, 2004).

(b) INK4 Family Members

INK4 proteins inhibit S-phase entry by preventing the formation of active cyclin D/CDK4 holoenzymes, due to the formation of binary complexes between the INK inhibitor and the catalytic subunit CDK4 (Quelle *et al.*, 1995, 1997).

The *Ink4* locus is composed of several genes identified as *Ink4a*, *Ink4b*, *Ink4c* and *Ink4d*. While each of the *Ink4 b-d* genes encodes for one protein named on the basis of the molecular weight p15INK4b, p18INK4c, p19INK4d, the *Ink4a* locus is unusual because its second exon contributes coding sequences to two distinct reading frames resulting in two proteins: p16INK4a and p19ARF (Zindy *et al.*, 1997).

In developing mouse embryos, only p18INK4c and p19INK4d have been identified (Zindy *et al.*, 1997). P18INK4c is preferentially localized in neurons as they exited from the cell cycle (Zindy *et al.*, 1999), whereas p19INK4d is mainly detected in post-mitotic neurons and expressed at high levels in the adult brain (Zindy *et al.*, 1999), often together with p27Kip1 (vanLookeren-Campagne and Gill, 1998). In the neonatal rat SVZ, p19INK4d levels are low in proliferating cells at the anterior border of the SVZ and progressively increase in the migratory cells of the rostral migratory stream (Luskin and Coskun, 2002), thus suggesting that this molecule plays a critical role in the induction of cell cycle exit once the migrating cells have reached their final destination (Zindy *et al.*, 1999). Consistent with this interpretation, studies on mice with targeted deletion of two major cell cycle inhibitors, p18INK4c and p27Kip1 continue to proliferate even after the migratory period (Zindy *et al.*, 1997).

The results regarding the expression of p16INK4a and its possible role in cell cycle regulation of developing CNS are more controversial. While Northern and Western Blot analysis of extracts from developing mouse embryos (van Lookeren Campagne and Gill 1998) have not detected any p16INK4a signal in the brain, different results have been obtained in the developing rat, where p16INK4a is expressed at high levels in the proliferating cells of the VZ from E16 to E20 (Zindy *et al.*, 1997). Although p16Ink4a expression is apparently down-regulated in the rat brain also, there appears to be a general consensus on the increasing levels of this protein with increasing age of the animal (Zindy *et al.*, 1997). It is important to mention, however, that p16INK4a can be easily detected *in vitro*, in dissociated primary cultures, thus suggesting that the stress of culturing

could induce the expression of molecules that may not be present in an *in vivo* context (Jacobs *et al.*, 1999a, 1999b). As previously mentioned, the p16INK4a represents the alpha transcript of the *Ink4a* locus and represents an inhibitor of cyclin D/CDK complexes acting on pRb-E2F complexes. The other transcript of the same *Ink4a* locus is p19ARF (beta transcript) and it originates from a promoter some 15 kb upstream of the alpha transcript resulting in a different reading frame of exon 2 than the alpha transcript (see Fig. 2.6).

As a consequence, the beta transcript encodes a protein that has no sequence homology with p16INK4a and that activates p53 rather than the pRb pathway (Fig. 2.6). Given the importance of the *Ink4a* locus in the transcription of regulatory components for two growth-inhibitory pathways, Rb and p53, it becomes easier to understand the high incidence of deletions or inactivations observed in this locus in patients with brain tumors.

The INK4a proteins have not been detected in the developing SVZ, although presumably their expression increases with age. Given the importance of these molecules as modulators of the cell cycle, it becomes critical to understand the mechanisms regulating their expression. In this respect, it has been shown that a member of the polycomb family of chromatin modifiers called Bmi is expressed in the adult SVZ and acts as a potent repressor of the *Ink4a* locus (Molofsky *et al.*, 2003). Mice with targeted deletions in the Bmi gene have a significant decrease in proliferation of neonatal and adult SVZ cells together with a 20 fold induction of p16INK4a gene product and a 3 fold increase of p19ARF (Molofsky *et al.*, 2003). Besides proliferation, the increased levels of p16INK4a also modulate the ability of the stem cells to self-renew, thus supporting the importance of the *Ink4* locus as tumor suppressor.

Remarkably, however, spontaneous glial tumors are not observed in the *Ink4a/Arf* null mutants. Even though both GFAP⁺ astrocytes and nestin⁺ cells in these mice have the characteristics of “immortal” cells (Holland *et al.*, 1998a), they still require the delivery of a constitutively active form of the




FIGURE 2.6. **The INK4a locus.** The INK4a locus can generate two transcripts: p19 ARF that regulates p53 function and p16INK4a, that modulates the activity of CDK4 and therefore regulates the Rb pathway.

EGFR (Bachoo *et al.*, 2002) or of the activated forms of Ras or Akt (Uhrbom *et al.* 2002; Kamijo *et al.*, 1997) for neoplastic transformation. Finally, it is worth mentioning that mice with selective deletion of p19ARF, with intact p16INK4a, develop spontaneous gliomas (Sherr and Roberts 1995), thus arguing that p19ARF rather than p16INK4a is involved in the neoplastic transformation of SVZ cells.

(c) Kip Family Members

Inhibitors of the Kip family can bind CDK4/cylinD complexes, although with lower affinity than the INK4 proteins, but this event does not result in efficient functional inhibition of enzymatic activity (Polyak *et al.*, 1994; Toyoshima and Hunter, 1994; Sherr and Roberts, 1999). The ability of the Kips to inhibit S-phase entry is mediated by the formation of ternary complexes with cyclin A or E and CDK2 (Russo *et al.*, 1996). The inhibitory effect of the Kip molecules on cyclin/CDK complexes is two-fold: they prevent substrate binding and rearrange the amino-terminal lobe of CDK2; thus blocking ATP binding (Russo *et al.*, 1996). Three Kip inhibitors have been identified: p27Kip1 (Matsuoka *et al.*, 1995), p21Cip1/Waf1 and p57Kip2 (Sherr and Roberts 1999). The p57Kip2 inhibitor is found in the VZ and SVZ of the developing rat brain at E16 and E18, and higher levels of expression are observed in post-mitotic cells at E20 (van Lookeren Campagne and Gill, 1998). The p21Cip/Waf1 inhibitor is also detected at E16 and E18, but its expression is confined to the ependymal layer of the ventricle and the choroid plexus and dramatically decreases to undetectable levels in the adult brain (van Lookeren Campagne and Gill, 1998). In agreement with this expression pattern *p21Cip1*^{-/-} mice do not show any change in the proliferative ability of cells in the developing or mature brain in physiological conditions (Qiu *et al.*, 2004).

Of the three members of the Kip family, p27Kip1 is undoubtedly the most interesting. Its expression is detected in proliferating cells of the VZ at mid-gestation (van Lookeren Campagne and Gill, 1998) and its levels progressively increase with increasing numbers of cell divisions (Delalle *et al.*, 1999). Given the characteristic pattern of expression during the embryonic neuro-genetic period, it has been suggested that p27Kip1 accumulation is part of the mechanism regulating progressive lengthening of the cell cycle and/or increased probability of cell cycle exit (Tarui *et al.*, 2005). Studies on p27Kip1 ^{-/-} mice, however, have shown that the length of the cell cycle (T_c) of cortical embryonic progenitors is not affected by p27Kip1 loss of function (Goto *et al.*, 2004), although there is a definite increase in the probability of the cells to re-enter the cell cycle, and thus an increase of the proliferating population.

The expression of p27Kip1, however, persists in cells of the adult SVZ and in the rostral migratory stream, thus suggesting a role for this molecule also in the regulation of the proliferating population in the adult brain (van Lookeren

Campagne and Gill, 1998). Mice with targeted deletions in the first exon of p27Kip1 show a selective increase in the number of transit amplifying progenitors concomitant with a reduction in the number of neuroblasts and no change in the number of stem cells (Doetsch *et al.*, 2002b). This indicates that cell cycle regulation of SVZ adult progenitors is remarkably cell-type specific with p27Kip1 being a key regulator of cell division in transit amplifying progenitors, but not of the slow proliferating stem cells (Doetsch *et al.*, 2002b). In vitro studies on neurospheres cultured from the neonatal SVZ support this interpretation. The levels of p27Kip1 are low in proliferating neurospheres, they increase during the early stages of differentiation and decrease again with time, in culture, thus indicating a possible role for this protein in regulating the cell cycle of immature, but not stem cells or the more mature neuroblasts (Jori *et al.*, 2003). Together, these data suggest that distinct molecular pathways may be activated in physiological and pathological conditions in order to modulate the number of neural stem cells (Fig. 2.7).

(d) p53 Pathway

The tumor-suppressor gene p53 is an important checkpoint for mammalian cells in the G1 phase of the cell cycle. Upon genotoxic stress, irradiation, DNA damage, oxidative stress or glucose deprivation, this molecule activates a transcriptional response resulting in either exit from the cell cycle (possibly mediated by up-regulation of p21Cip1/Waf1) or apoptosis. In the developing brain, however, p53 expression is most abundant in proliferating




FIGURE 2.7. Schematic representation of extracellular signals and intracellular molecules regulating the decision of a cell in the G1 phase of the cell cycle. Although it is not clear whether mitogenic and anti-mitogenic signals affect the same cellular effector molecules in SVZ cells, it is likely that activation of active cyclin/CDK complexes result in proliferation, while their inhibition by Kip and INK family members may result in cell cycle exit.

cell populations of the embryonic and postnatal rat brain, and is not observed in regions undergoing spontaneous apoptosis (Donehower *et al.*, 1992). At E14, its levels are very high in proliferating cells of the ventricular zone, while from E16 to E20, it is also expressed in the SVZ and the cortical plate. The expression of p53 decreases postnatally, but it remains quite high in the postnatal rostral migratory stream and in the subventricular zone, where it persists together with p27Kip1 (van Lookeren Campagne and Gill, 1998). Interestingly, the pattern of expression of p21Cip1/Waf1, one of the downstream transcriptional targets of p53, is quite different, indicating that the role of p53 in cell cycle regulation of adult neural stem cells is independent of p21Cip1/Waf1 expression. Despite the high levels of p53 detected in the VZ and SVZ of the developing rat brain, p53 null mice develop normally, and do not display any major defects in brain histoarchitecture (Donehower *et al.*, 1992). Intriguingly, however, they do display increased susceptibility to the development of glial tumors after transplacental exposure to mutagens (Leonard *et al.*, 2001). Current studies in our laboratory support the hypothesis that the increased susceptibility of these mice to brain tumors is secondary to a specific role of this molecule in modulating the number of adult neural stem cells in vivo (SGP and PCB unpublished).

In vitro, the levels of the cell cycle regulator p53 are quite low in proliferating neurospheres generated from neonatal rats and maintained in EGF and its transcript levels are significantly higher in cells differentiated after mitogen withdrawal (Nakamura *et al.*, 2000; Jori *et al.*, 2003). Higher p53 levels correlate with increased apoptotic index in vitro after 3-7 days in culture. Increased protein levels, however, are observed only after 21 days in differentiating conditions, and correlate with the detection of high levels of neuronal and glial markers, thus suggesting a dual role for this molecule in apoptosis and in differentiation or lineage commitment of neural stem cells.

Other Intracellular Signaling Molecules

Emx2

Emx2 and the related gene *Emx1* are the vertebrate homologues of the *Drosophila* gene *Empty spiracles (ems)* involved in cephalic development (Mallamaci *et al.*, 1998). The distinct expression pattern during late embryonic development with *Emx2* expression restricted to the VZ, and *Emx1* strongly expressed in the subplate and cortical plate (Gulisano *et al.*, 1996), suggests that these two transcription factors play distinct roles in the developing nervous system. *Emx2* is involved in proliferation and migration while *Emx1* seems to affect neurogenesis (Yoshida *et al.*, 1997). *Emx1* null mice, however, do not have the corpus callosum and show only subtle defects in cerebral cortex (Pellegrini *et al.*, 1996), while *Emx2* null mice display major alterations of the brain histoarchitecture (Mallamaci *et al.*, 2000; Tole *et al.*, 2000). Recent studies on *Emx2* null mice have shown significant enlargement of the proliferative ventricular and subventricular zones (Galli *et al.*, 2002),

thus suggesting that this molecule acts as a negative regulator of proliferation of neural precursors and adult neural stem cells. *Emx2* is expressed *in vivo* in the adult SVZ (Gangemi *et al.*, 2001; Galli *et al.*, 2002) and in the rostral migratory stream, and *in vitro* in multipotent neural precursors (Galli *et al.*, 2002). Its expression is significantly decreased when these stem cells differentiate into neurons and glia (Gangemi *et al.*, 2001; Galli *et al.*, 2002). Gain-of-function studies by over-expressing *Emx2* decrease the proliferative rate of cells while retaining their differentiative potential (Gangemi *et al.*, 2001; Galli *et al.*, 2002). Based on these *in vivo* and *in vitro* studies, it can be concluded that *Emx2* acts as a negative regulator of proliferation of adult neural stem cells.

Vax 1

The homeobox *Vax1* is a homologue of *Emx2* and is also strongly expressed in the embryonic and adult SVZ and in the RMS (Soria *et al.*, 2004). In the absence of *Vax1*, embryonic precursor cells proliferate 100 times more than wild-type controls, *in vitro*. In addition, the SVZ of *Vax1* null mice shows signs of hyperplasia and disorganization (Soria *et al.*, 2004). Together, these data suggested that, like *Emx2*, the transcription factor *Vax1* is an important regulator of proliferation of SVZ cells.

PTEN

PTEN is a lipid phosphatase originally cloned as a tumor suppressor for glioma (Li *et al.*, 1997; Tamura *et al.*, 1998; Datta *et al.*, 1999). PTEN is a phosphatidylinositol (PIP) phosphatase, responsible for the dephosphorylation of PIP3, thus antagonizing the role of the survival kinases PI3K and Akt and rendering the cells more susceptible to apoptosis (Groszer *et al.*, 2001). In addition, PTEN is responsible for the dephosphorylation of the focal adhesion kinase FAK, resulting in the inhibition of cell migration (Groszer *et al.*, 2001). In the adult brain, PTEN is expressed mainly in neurons and is found both in the nucleus and cytoplasm of cells in the olfactory bulb, in the SVZ and in large projection neurons. Given the importance of this signaling molecule in regulating multiple pathways, several groups have generated conditional knockout mice using the Cre-lox system. The first to be reported is the PTEN deleted by Cre expression in nestin+ cells (Backman *et al.*, 2001; Kwon *et al.*, 2001). These mice show increased proliferation and decreased apoptosis of cells lining the ventricular walls with a dramatic brain enlargement and death immediately after birth (Li *et al.*, 2003). Very different is the phenotype of mice where PTEN is deleted in cells expressing Cre from the GFAP promoter (Recht *et al.*, 2003; Berger *et al.*, 2004). In this case, no change in proliferation or apoptosis has been reported, although the mice displayed an abnormal organization of the cerebellum. These data clearly indicate that the effect of PTEN is cell-context dependent and is affected by the intracellular and extracellular milieu, possibly due to the cross-talk with distinct signaling pathways that are active in different cells at different times.

Conclusions

Although stem cell therapy has been proposed for therapeutic strategies aimed at repairing functions, it is important to realize that as yet, relatively little is known about the behavior of embryonic and adult stem cells in terms of responsiveness to extracellular cues and intracellular signaling molecules.

The challenge that awaits ahead is to define possible differences in intracellular signaling molecules between embryonic and adult derived neural stem cells that may underlie the distinctive responsiveness of these different cell types to external signals. A better understanding of the mechanisms regulating proliferation and differentiation of multipotent progenitors into differentiated neurons, astrocyte and oligodendrocytes is, therefore, essential for developing a realistic frame of therapeutic intervention while preventing undesirable - and yet possible-neoplastic transformation of adult neural stem cells.

Acknowledgments. The authors are grateful to Dr Aixiao Liu and Siming Shen for critical reading of the text; to Dr Richard Nowakowski and Dr Charles French-Constant for valuable comments and to Ms Bonnefil Valentina for constant support. Dr Casaccia-Bonnefil is supported by funds from NIH-NINDS and from the National Multiple Sclerosis Society. Dr Gil-Perotin is supported by a fellowship from Instituto Salud Carlos III.

References

- Ahmed, S., Reynolds, B.A. and Weiss, S. (1995). BDNF enhances the differentiation but not the survival of CNS stem cell-derived neuronal precursors. *J. Neurosci.* 15(8): 5765–5778.
- Aloe, L. (2004). Rita Levi-Montalcini: The discovery of nerve growth factor and modern neurobiology. *Trends Cell. Biol.* 14(7): 395–399.
- Alvarez-Buylla, A. and Lim, D.A. (2004). For the long run: Maintaining germinal niches in the adult brain. *Neuron* 41(5): 683–686.
- Anderson, K.D., Alderson, R.F., Altar, C.A., DiStefano, P.S., Corcoran, T.L., Lindsay, R.M. and Wiegand, S.J. (1995). Differential distribution of exogenous BDNF, NGF, and NT-3 in the brain corresponds to the relative abundance and distribution of high-affinity and low-affinity neurotrophin receptors. *J. Comp. Neurol.* 357(2): 296–317.
- Anderson, K.D., Panayotatos, N., Corcoran, T.L., Lindsay, R.M. and Wiegand, S.J. (1996). Ciliary neurotrophic factor protects striatal output neurons in an animal model of Huntington disease. *Proc. Natl. Acad. Sci. U. S. A.* 93(14): 7346–7351.
- Anton, E.S., Ghashghaei, H.T., Weber, J.L., McCann, C., Fischer, T.M., Cheung, I.D., Gassmann, M., Messing, A., Klein, R., Schwab, M.H., Lloyd, K.C. and Lai, C. (2004). Receptor tyrosine kinase ErbB4 modulates neuroblast migration and placement in the adult forebrain. *Nat. Neurosci.* 7(12): 1319–1328.

- Arsenijevic, Y. and Weiss, S. (1998). Insulin-like growth factor-I is a differentiation factor for postmitotic CNS stem cell-derived neuronal precursors: Distinct actions from those of brain-derived neurotrophic factor. *Neurosci. J.* 18(6): 2118–2128.
- Arsenijevic, Y., Weiss, S., Schneider, B. And Aebischer, P. (2001). Insulin-like growth factor-I is necessary for neural stem cell proliferation and demonstrates distinct actions of epidermal growth factor and fibroblast growth factor-2. *Neurosci J.* 21(18): 7194–7202.
- Artavanis-Tsakonas, S., Matsuno, K. and Fortini, M.E. (1995). Notch. signaling. *Science.* 268(5208): 225–232.
- Baas, D., Bourbeau, D., Sarlieve, L.L., Ittel, M.E., Dussault, J.H. and Puymirat, J. (1997). Oligodendrocyte maturation and progenitor cell proliferation are independently regulated by thyroid hormone. *Glia* 19(4): 324–332.
- Bachoo RM., Maher EL, Ligon KL, Sharpless NE, Chan SS, You MJ, Tang Y, DeFrances J, Stover E, Weissleder R, Rowitch DH, Louis DN, DePinho RA. (2002). Epidermal growth factor receptor and Ink4a/Arf: Convergent mechanisms governing terminal differentiation and transformation along the neural stem cell to astrocyte axis. *Cancer Cell.* (2002) Apr. 1(3):269–277.
- Backman, S.A., Stambolic, V., Suzuki, A., Haight, J., Elia, A., Pretorius, J., Tsao, M.S., Shannon, P., Bolon, B., Ivy G.O., and Mak, T.W. (2001). Deletion of Pten in mouse brain causes seizures, ataxia and defects in soma size resembling Lhermitte-Duclos disease. *Nat. Genet.* 29(4): 396–403.
- Baker, J., Liu, J.P., Robertson, E.J. and Efstratiadis, A. (1993). Role of insulin-like growth factors in embryonic and postnatal growth. *Cell.* 75(1): 73–82.
- Banasr, M., Hery, M., Printemps, R. and Daszuta, A. (2004). Serotonin-induced increases in adult cell proliferation and neurogenesis are mediated through different and common 5-HT receptor subtypes in the dentate gyrus and the subventricular zone. *Neuropsychopharmacology* 29(3): 450–460.
- Barker, P.A. (2004). p75^{NTR} is positively promiscuous: Novel partners and new insights. *Neuron* 42(4): 529–533.
- Barres, B.A., Burne, J.F., Holtmann, B., Thoenen, H., Sendtner, M. and Raff, M.C. (1996). Ciliary neurotrophic factor enhances the rate of oligodendrocyte generation. *Mol. Cell. Neurosci.* 8(2-3): 146–156.
- Baumann, N. and Pham-Dinh, D. (2001). Biology of oligodendrocyte and myelin in the mammalian central nervous system. *Physiol. Rev.* 81(2): 871–927.
- Bayer, S.A. and Altman, J. (1991). Development of the endopiriform nucleus and the claustrum in the rat brain. *Neuroscience* 45(2): 391–412.
- Bazan, J.F. (1991). Neuropoietic cytokines in the hematopoietic fold. *Neuron* 7(2): 197–208.
- Beck, K.D., Powell-Braxton, L., Widmer, H.R., Valverde, J. and Hefti, F. (1995). Igf1 gene disruption results in reduced brain size, CNS hypomyelination, and loss of hippocampal granule and striatal parvalbumin-containing neurons. *Neuron* 14(4): 717–730.
- Beijersbergen, R.L., Kerkhoven, R.M., Zhu, L., Carlee, L., Voorhoeve P.M. and Bernards, R. (1994). E2F-4, a new member of the E2F gene family, has oncogenic activity and associates with p107 in vivo. *Genes Dev.* 8(22): 2680–2690.
- Benoit, B.O., Savarese, T., Joly, M., Engstrom, C.M., Pang, L., Reilly, J., Recht, L.D., Ross, A.H. and Quesenberry, P.J. (2001). Neurotrophin channeling of neural progenitor cell differentiation. *J. Neurobiol.* 46(4): 265–280.

- Berger, F., Gay, E., Pelletier, L., Tropel P., and Wion, D. (2004). Development of gliomas: Potential role of asymmetrical cell division of neural stem cells. *Lancet. Oncol.* 5(8): 511–514.
- Berkemeier, L.R., Winslow, J.W., Kaplan, D.R., Nikolics, K., Goeddel, D.V. and Rosenthal, A. (1991). Neurotrophin-5: A novel neurotrophic factor that activates trk and trkB. *Neuron* 7(5): 857–866.
- Bondy, C.A., Werner, H., Roberts, Jr., C.T. and LeRoith, D. (1990). Cellular pattern of insulin-like growth factor-I (IGF-I) and type I IGF receptor gene expression in early organogenesis: Comparison with IGF-II gene expression. *Mol. Endocrinol.* 4(9): 1386–1398.
- Bonni, A., Sun, Y., Nadal-Vicens, M., Bhatt, A., Frank, D.A., Rozovsky, I., Stahl, N., Yancopoulos, G.D. and Greenberg, M.E. (1997). Regulation of gliogenesis in the central nervous system by the JAK-STAT signaling pathway. *Science* 278(5337): 477–483.
- Bradley, D.J., Towle H.C. and Young, 3rd (1992). Spatial and temporal expression of alpha-and beta-thyroid hormone receptor mRNAs, including the beta 2-subtype, in the developing mammalian nervous system. *J. Neurosci.* 12(6): (2288–2302).
- Brannvall, K., Korhonen L. and Lindholm, D. (2002). Estrogen-receptor-dependent regulation of neural stem cell proliferation and differentiation. *Mol Cell Neurosci.* 21(3): 512–520.
- Brehm, A., Miska, E.A., McCance, D.J., Reid, J.L., Bannister A. J. and Kouzarides, T. (1998). Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature* 391(6667): 597–601.
- Breier, G., Albrecht, U., Sterrer, S. and Risau, W. (1992). Expression of vascular endothelial growth factor during embryonic angiogenesis and endothelial cell differentiation. *Development* 114(2): 521–532.
- Bridges, R.S. and Grattan, D.R. (2003). Prolactin-induced neurogenesis in the maternal brain. *Trends Endocrinol. Metab.* 14(5): 199–201.
- Buemi, M., Cavallaro, E., Floccari, F., Sturiale, A., Aloisi, C., Trimarchi, M., Grasso, G., Corica F. and Frisina, N., (2002). Erythropoietin and the brain: From neurodevelopment to neuroprotection. *Clin. Sci. (Lond.)* 103(3): 275–282.
- Burke, D., Wilkes, D., Blundell T. L. and Malcolm, S. (1998). Fibroblast growth factor receptors: Lessons from the genes. *Trends Biochem. Sci.* 23(2): 59–62.
- Burrows, R.C., Wancio, D., Levitt, P. and Lillien, L. (1997). Response diversity and the timing of progenitor cell maturation are regulated by developmental changes in EGFR expression in the cortex. *Neuron* 19(2): 251–267.
- Caille, I., Allinquant, B., Dupont, E., Bouillot, C., Langer, A., Muller, U. and Prochiantz, A. (2004). Soluble form of amyloid precursor protein regulates proliferation of progenitors in the adult subventricular zone. *Development* 131(9): 2173–2181.
- Calaora, V., Rogister, B., Bismuth, K., Murray, K., Brandt, H., Leprince, P., Marchionni, M. and Dubois-Dalcq, M. (2001). Neuregulin signaling regulates neural precursor growth and the generation of oligodendrocytes in vitro. *J Neurosci* 21(13): 4740–4751.
- Callaghan, D.A., Dong, L., Callaghan, S.M., Hou, Y.X., Dagnino, L. and Slack, R.S. (1999). Neural precursor cells differentiating in the absence of Rb exhibit delayed terminal mitosis and deregulated E2F 1 and 3 activity. *Dev Biol* 207(2): 257–270.
- Carnot, P. and Deflandre, C. (1906). Sur l'activité hémopoïétique des différents organes au cours de la régénération du sang. *C.R. Acad. Sci. Paris* (143): 432–435.

- Carson, M.J., Behringer, R.R., Brinster, R.L. and McMorris, F.A. (1993). Insulin-like growth factor I increases brain growth and central nervous system myelination in transgenic mice. *Neuron* 10(4): 729–740.
- Caviness, V.S., Jr., Takahashi, T. and Nowakowski, R.S. (1995). Numbers, time and neocortical neurogenesis: A general developmental and evolutionary model. *Trends Neurosci* 18(9): 379–383.
- Chambers, C.B., Peng, Y., Nguyen, H., Gaiano, N., Fishell, G. and Nye, J.S. (2001). Spatiotemporal selectivity of response to Notch1 signals in mammalian forebrain precursors. *Development* 128(5): 689–702.
- Chao, M.V. (2003). Neurotrophins and their receptors: A convergence point for many signalling pathways. *Nat. Rev. Neurosci.* 4(4): 299–309.
- Chenn, A. and Walsh, C.A. (2002). Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. *Science* 297(5580): 365–369.
- Chiang, C., Litingtung, Y., Lee, E., Young, K.E., Corden, J.L., Westphal, H. and Beachy, P.A. (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* 383(6599): 407–413.
- Chipperfield, J.G., Perry, R.P. and Weiner, B. (2003). Discrete emotions in later life. *J Gerontol B Psychol. Sci. Soc. Sci.* 58(1): P23–P34.
- Chojnacki, A., Shimazaki, T., Gregg, C., Weinmaster, G. and Weiss, S. (2003). Glycoprotein 130 signaling regulates Notch1 expression and activation in the self-renewal of mammalian forebrain neural stem cells. *J. Neurosci.* 23(5): 1730–1741.
- Ciccolini, F. and Svendsen, C.N. (1998). Fibroblast growth factor 2 (FGF-2) promotes acquisition of epidermal growth factor (EGF) responsiveness in mouse striatal precursor cells: Identification of neural precursors responding to both EGF and FGF-2. *J. Neurosci.* 18(19): 7869–7880.
- Clarke, A.R., Maandag, E.R., van Roon, M., van der Lugt, N.M., van der Valk, M., Hooper, M.L., Berns, A. and te Riele, H. (1992). Requirement for a functional Rb-1 gene in murine development. *Nature* 359(6393): 328–330.
- Cobrinik, D., Lee, M.H., Hannon, G., Mulligan, G., Bronson, R.T., Dyson, N., Harlow, E., Beach, D., Weinberg, R.A. and Jacks, T. (1996). Shared role of the pRB-related p130 and p107 proteins in limb development. *Genes Dev.* 10(13): 1633–1644.
- Conover, J.C. and Allen, R.L. (2002). The subventricular zone: New molecular and cellular developments. *Cell Mol. Life Sci.* 59(12): 2128–2135.
- Conover, J.C., Doetsch, F., Garcia-Verdugo, J.M., Gale, N.W., Yancopoulos, G.D. and Alvarez-Buylla, A. (2000). Disruption of Eph/ephrin signaling affects migration and proliferation in the adult subventricular zone. *Nat. Neurosci.* 3(11): 1091–1107.
- Conover, J.C., Ip, N.Y., Poueymirou, W.T., Bates, B., Goldfarb, M.P., DeChiara, M.T. and Yancopoulos, G.D. (1993). Ciliary neurotrophic factor maintains the pluripotentiality of embryonic stem cells. *Development* 119(3): 559–565.
- Conrad, C.D., Grote, K.A., Hobbs, R.J. and Ferayorni, A. (2003). Sex differences in spatial and non-spatial Y-maze performance after chronic stress. *Neurobiol. Learn. Mem.* 79(1): 32–40.
- Constam, D.B., Schmid, P., Aguzzi, A., Schachner, M. and Fontana, A. (1994). Transient production of TGF-beta 2 by postnatal cerebellar neurons and its effect on neuroblast proliferation. *Eur. J. Neurosci.* 6(5): 766–778.
- Cooper, O. and Isacson, O. (2004). Intra-striatal transforming growth factor alpha delivery to a model of Parkinson's disease induces proliferation and migration of

- endogenous adult neural progenitor cells without differentiation into dopaminergic neurons. *J. Neurosci.* 24(41): 8924–8931.
- Corfas, G., Rosen, K.M., Aratake, H., Krauss, R. and Fischbach, G.D. (1995). Differential expression of ARIA isoforms in the rat brain. *Neuron* 14(1): 103–115.
- Coronas, V., Bantubungi, K., Fombonne, J., Krantic, S., Schiffmann, S.N. and Roger, M. (2004). Dopamine D3 receptor stimulation promotes the proliferation of cells derived from the post-natal subventricular zone. *J. Neurochem.* 91(6): 1292–1301.
- Coskun, V. and Luskin, M.B. (2001). The expression pattern of the cell cycle inhibitor p19(INK4d) by progenitor cells of the rat embryonic telencephalon and neonatal anterior subventricular zone. *J. Neurosci.* 21(9): 3092–3103.
- Craig, C.G., Tropepe, V., Morshead, C.M., Reynolds, B.A., Weiss, S. and van der Kooy, D. (1996). In vivo growth factor expansion of endogenous subependymal neural precursor cell populations in the adult mouse brain. *J. Neurosci.* 16(8): 2649–2658.
- Dahmane, N. and Ruiz-i-Altaba, A. (1999). Sonic hedgehog regulates the growth and patterning of the cerebellum. *Development* 126(14): 3089–3100.
- Dallner, C., Woods, A.G., Deller, T., Kirsch, M. and Hofmann, H.D. (2002). CNTF and CNTF receptor alpha are constitutively expressed by astrocytes in the mouse brain. *Glia.* 37(4): 374–408.
- Datta, S.R., Brunet, A. and Greenberg, M.E. (1999). Cellular survival: A play in three Akts. *Genes Dev.* 13(22): 2905–2927.
- Davis, S., Aldrich, T.H., Stahl, N., Pan, L., Taga, T., Kishimoto, T., Ip, N.Y. and Yancopoulos, G.D. (1993). LIFR beta and gp130 as heterodimerizing signal transducers of the tripartite CNTF receptor. *Science* 260(5115): 1805–1808.
- De Felici, M. and Dolci, S. (1991). Leukemia inhibitory factor sustains the survival of mouse primordial germ cells cultured on TM4 feeder layers. *Dev. Biol.* 147(1): 281–284.
- DeChiara, T.M., Vejsada, R., Poueymirou, W.T., Acheson, A., Suri, C., Conover, J.C., Friedman, B., McClain, J., Pan, L., Stahl, N. and et al. (1995). Mice lacking the CNTF receptor, unlike mice lacking CNTF, exhibit profound motor neuron deficits at birth. *Cell* 83(2): 313–322.
- Delalle, I., Takahashi, T., Nowakowski, R.S., Tsai, L.H. and Caviness, Jr., V.S. (1999). Cyclin E-p27 opposition and regulation of the G1 phase of the cell cycle in the murine neocortical PVE: A quantitative analysis of mRNA in situ hybridization. *Cereb Cortex* 9(8): 824–832.
- Deng, C., Bedford, M., Li, C., Xu, X., Yang, X., Dunmore, J. and Leder, P. (1997). Fibroblast growth factor receptor-1 (FGFR-1) is essential for normal neural tube and limb development. *Dev. Biol.* 185(1): 42–54.
- Deng, C.X., Wynshaw-Boris, A., Shen, M.M., Daugherty, C., Ornitz, D.M. and Leder, P. (1994). Murine FGFR-1 is required for early postimplantation growth and axial organization. *Genes Dev.* 8(24): 3045–3057.
- Dhawan, B.N., Cesselin, F., Raghbir, R., Reisine, T., Bradley, P.B., Portoghese, P.S. and Hamon, M. (1996). International Union of Pharmacology. XII. Classification of opioid receptors. *Pharmacol. Rev.* 48(4): 567–592.
- Diaz, J., Ridray, S., Mignon, V., Griffon, N., Schwartz, J.C. and Sokoloff, P. (1997). Selective expression of dopamine D3 receptor mRNA in proliferative zones during embryonic development of the rat brain. *J. Neurosci.* 17(11): 4282–4292.
- Doetsch, F. (2003). A niche for adult neural stem cells. *Curr. Opin. Genet. Dev.* 13(5): 543–550.

- Doetsch, F., Garcia-Verdugo, J.M. and Alvarez-Buylla, A. (1997). Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain. *J. Neurosci.* 17(13): 5046–5061.
- Doetsch, F., Petreanu, L., Caille, I., Garcia-Verdugo, J.M. and Alvarez-Buylla, A. (2002a). EGF converts transit-amplifying neurogenic precursors in the adult brain into multipotent stem cells. *Neuron* 36(6): 1021–1034.
- Doetsch, F., Verdugo, J.M., Caille, I., Alvarez-Buylla, A., Chao, M.V. and Casaccia-Bonnel, P. (2002b). Lack of the cell-cycle inhibitor p27Kip1 results in selective increase of transit-amplifying cells for adult neurogenesis. *J. Neurosci.* 22(6): 2255–2264.
- Donehower, L.A., Harvey, M., Slagle, B.L., McArthur, M.J., Montgomery, Jr., C.A., Butel, J.S. and Bradley, A. (1992). Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* 356(6366): 215–221.
- Drago, J., Murphy, M., Carroll, S.M., Harvey, R.P. and Bartlett, P.F. (1991). Fibroblast growth factor-mediated proliferation of central nervous system precursors depends on endogenous production of insulin-like growth factor I. *Proc. Natl. Acad. Sci. U. S. A.* 88(6): 2199–2203.
- Emerich, D.F., Lindner, M.D., Winn, S.R., Chen, E.Y., Frydel, B.R. and Kordower, J.H. (1996). Implants of encapsulated human CNTF-producing fibroblasts prevent behavioral deficits and striatal degeneration in a rodent model of Huntington's disease. *J. Neurosci.* 16(16): 5168–5181.
- Ericson, J., Morton, S., Kawakami, A., Roelink, H. and Jessell, T.M. (1996). Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity. *Cell* 87(4): 661–673.
- Ernsberger, U., Sendtner, M. and Rohrer, H. (1989). Proliferation and differentiation of embryonic chick sympathetic neurons: Effects of ciliary neurotrophic factor. *Neuron* 2(3): 1275–1284.
- Ezzeddine, Z.D., Yang, X., DeChiara, T., Yancopoulos, G. and Cepko, C.L. (1997). Postmitotic cells fated to become rod photoreceptors can be respecified by CNTF treatment of the retina. *Development* 124(5): 1055–1067.
- Ferguson, K.L. and Slack, R.S. (2001). The Rb pathway in neurogenesis. *Neuroreport* 12(9): A55–A62.
- Fernandez, M., Pirondi, S., Manservigi, M., Giardino, L. and Calza, L. (2004). Thyroid hormone participates in the regulation of neural stem cells and oligodendrocyte precursor cells in the central nervous system of adult rat. *Eur. J. Neurosci.* 20(8): 2059–2070.
- Ferrara, N., Carver-Moore, K., Chen, H., Dowd, M., Lu, L., O'Shea, K.S., Powell-Braxton, L., Hillan, K.J. and Moore, M.W. (1996). Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature* 380(6573): 439–442.
- Fiore, M., Amendola, T., Triaca, V., Alleva, E. and Aloe, L. (2005). Fighting in the aged male mouse increases the expression of TrkA and TrkB in the subventricular zone and in the hippocampus. *Behav Brain Res.* 157(2): 351–362.
- Fisher, J.W. (2003). Erythropoietin: Physiology and pharmacology update. *Exp. Biol. Med.* (Maywood) 228(1): 1–14.
- Flanders, K.C., Ludecke, G., Engels, S., Cissel, D.S., Roberts, A.B., Kondaiah, P., Lafyatis, R., Sporn, M.B. and Unsicker, K. (1991). Localization and actions of transforming growth factor-beta s in the embryonic nervous system. *Development* 113(1): 183–191.

- Frye, C.A., Petralia, S.M. and Rhodes, M.E. (2000). Estrous cycle and sex differences in performance on anxiety tasks coincide with increases in hippocampal progesterone and 3 α ,5 α -THP. *Pharmacol. Biochem. Behav.* 67(3): 587–596.
- Gago, N., Avellana-Adalid, V., Evercooren, A.B. and Schumacher, M. (2003). Control of cell survival and proliferation of postnatal PSA-NCAM(+) progenitors. *Mol. Cell. Neurosci.* 22(2): 162–178.
- Gaiano, N., Nye, J.S. and Fishell, G. (2000). Radial glial identity is promoted by Notch1 signaling in the murine forebrain. *Neuron* 26(2): 395–404.
- Galli, R., Fiocco, R., De Filippis, L., Muzio, L., Gritti, A., Mercurio, S., Broccoli, V., Pellegrini, M., Mallamaci, A. and Vescovi, A.L. (2002). Emx2 regulates the proliferation of stem cells of the adult mammalian central nervous system. *Development* 129(7): 1633–1644.
- Galli, R., Pagano, S.F., Gritti, A. and Vescovi, A.L. (2000). Regulation of neuronal differentiation in human CNS stem cell progeny by leukemia inhibitory factor. *Dev. Neurosci.* 22(1-2): 86–95.
- Gangemi, R.M., Daga, A., Marubbi, D., Rosatto, N., Capra, M.C. and Corte, G. (2001). Emx2 in adult neural precursor cells. *Mech. Dev.* 109(2): 323–329.
- Garcia-Segura, L.M., Perez, J., Pons, S., Rejas, M.T. and Torres-Aleman, I. (1991). Localization of insulin-like growth factor I (IGF-I)-like immunoreactivity in the developing and adult rat brain. *Brain Res.* 560(1-2): 167–174.
- Garcia-Verdugo, J.M., Doetsch, F., Wichterle, H., Lim, D.A. and Alvarez-Buylla, A. (1998). Architecture and cell types of the adult subventricular zone: In search of the stem cells. *J. Neurobiol.* 36(2): 234–248.
- Garcion, E., Halilagic, A., Faissner, A. and French-Constant, C. (2004). Generation of an environmental niche for neural stem cell development by the extracellular matrix molecule tenascin C. *Development* 131(14): 3423–3432.
- Garofalo, R.S. and Rosen, O.M. (1988). Tissue localization of *Drosophila melanogaster* insulin receptor transcripts during development. *Mol. Cell. Biol.* 8(4): 1638–1647.
- Gascon, E., Vutskits, L., Zhang, H., Barral-Moran, M.J., Kiss, P.J., Mas, C. and Kiss, J.Z. (2005). Sequential activation of p75 and TrkB is involved in dendritic development of subventricular zone-derived neuronal progenitors in vitro. *Eur. J. Neurosci.* 21(1): 69–80.
- Gaspar, P., Cases, O. and Maroteaux, L. (2003). The developmental role of serotonin: News from mouse molecular genetics. *Nat. Rev. Neurosci.* 4(12): 1002–1012.
- Gearing, D.P. and Bruce, A.G. (1992). Oncostatin M binds the high-affinity leukemia inhibitory factor receptor. *New. Biol.* 4(1): 61–65.
- Gearing, D.P., Comeau, M.R., Friend, D.J., Gimpel, S.D., Thut, C.J., McGourty, J., Brasher, K.K., King, J.A., Gillis, S., Mosley, B. and *et al.* (1992). The IL-6 signal transducer, gp130: An oncostatin M receptor and affinity converter for the LIF receptor. *Science* 255(5050): 1434–1437.
- Gearing, D.P., Thut, C.J., VandeBos, T., Gimpel, S.D., Delaney, P.B., King, J., Price, V., Cosman, D. and Beckmann, M.P. (1991). Leukemia inhibitory factor receptor is structurally related to the IL-6 signal transducer, gp130. *Embo. J.* 10(10): 2839–2848.
- Giardino, L., Bettelli, C. and Calza, L. (2000). In vivo regulation of precursor cells in the subventricular zone of adult rat brain by thyroid hormone and retinoids. *Neurosci. Lett.* 295(1-2): 17–20.

- Gill, R.M., Slack, R., Kiess, M. and Hamel, P.A. (1998). Regulation of expression and activity of distinct pRB, E2F, D-type cyclin, and CKI family members during terminal differentiation of P19 cells. *Exp. Cell Res.* 244(1): 157–170.
- Ginsberg, D., Vairo, G., Chittenden, T., Xiao, Z.X., Xu, G., Wydner, K.L., DeCaprio, J.A., Lawrence, J.B. and Livingston, D. M. (1994). E2F-4, a new member of the E2F transcription factor family, interacts with p107. *Genes Dev.* 8(22): 2665–2679.
- Giuliani, A., D’Intino, G., Paradisi, M., Giardino, L. and Calza, L. (2004). p75(NTR)-Immunoreactivity in the subventricular zone of adult male rats: Expression by cycling cells. *J. Mol. Histol.* 35(8-9): 749–758.
- Goto, T, Mitsuhashi, T, Takahashi, T. (2004). Altered Patterns of Neuron Production in the p27 Knockout Mouse. *Dev. Neurosci.* 26:208–217.
- Gould, E. and Tanapat, P. (1997). Lesion-induced proliferation of neuronal progenitors in the dentate gyrus of the adult rat. *Neuroscience.* 80(2): 427–436.
- Graham, A., Koentges, G. and Lumsden, A. (1996). Neural crest apoptosis and the establishment of craniofacial pattern: An honorable death. *Mol. Cell Neurosci.* 8(2-3): 76–83.
- Grinspan, J.B., Edell, E., Carpio, D.F., Beesley, J.S., Lavy, L., Pleasure, D. and Golden, J.A. (2000). Stage-specific effects of bone morphogenetic proteins on the oligodendrocyte lineage. *J. Neurobiol.* 43(1): 1–17.
- Gritti, A., Frolichsthal-Schoeller, P., Galli, R., Parati, E.A., Cova, L., Pagano, S.F., Bjornson, C.R. and Vescovi, A.L. (1999). Epidermal and fibroblast growth factors behave as mitogenic regulators for a single multipotent stem cell-like population from the subventricular region of the adult mouse forebrain. *J. Neurosci.* 19(9): 3287–3297.
- Gross, R.E., Mehler, M.F., Mabie, P.C., Zang, Z., Santschi, L. and Kessler, J.A. (1996). Bone morphogenetic proteins promote astroglial lineage commitment by mammalian subventricular zone progenitor cells. *Neuron.* 17(4): 595–606.
- Groszer, M., Erickson, R., Scripture-Adams, D.D., Lesche, R., Trumpp, A., Zack, J.A., Kornblum, H.I. , Liu, X. and Wu, H. (2001). Negative regulation of neural stem/progenitor cell proliferation by the Pten tumor suppressor gene in vivo. *Science* 294(5549): 2186–2189.
- Gu, W., Brannstrom, T. and Wester, P. (2000). Cortical neurogenesis in adult rats after reversible photothrombotic stroke. *J. Cereb. Blood Flow Metab.* 20(8): 1166–1173.
- Guillin, O., Diaz, J., Carroll, P., Griffon, N., Schwartz, J.C. and Sokoloff, P. (2001). BDNF controls dopamine D3 receptor expression and triggers behavioural sensitization. *Nature* 411(6833): 86–89.
- Guillin, O., Griffon, N., Bezaud, E., Leriche, L., Diaz, J., Gross, C. and Sokoloff, P. (2003). Brain-derived neurotrophic factor controls dopamine D3 receptor expression: Therapeutic implications in Parkinson’s disease. *Eur. J. Pharmacol.* 480(1-3): 89–95.
- Guliano, M., Broccoli, V., Pardini, C. and Boncinelli, E. (1996). Emx1 and Emx2 show different patterns of expression during proliferation and differentiation of the developing cerebral cortex in the mouse. *Eur. J. Neurosci.* 8(5): 1037–1050.
- Hagg, T. and Varon, S. (1993). Ciliary neurotrophic factor prevents degeneration of adult rat substantia nigra dopaminergic neurons in vivo. *Proc. Natl. Acad. Sci. U. S. A.* 90(13): 6315–6319.
- Hall, A.C., Lucas, F.R. and Salinas, P.C. (2000). Axonal remodeling and synaptic differentiation in the cerebellum is regulated by WNT-7a signaling. *Cell* 100(5): 525–535.

- Hallbook, F., Ibanez, C.F. and Persson, H. (1991). Evolutionary studies of the nerve growth factor family reveal a novel member abundantly expressed in *Xenopus* ovary. *Neuron* 6(5): 845–858.
- Haughey, N.J., Liu, D., Nath, A., Borchard, A.C. and Mattson, M.P. (2002). Disruption of neurogenesis in the subventricular zone of adult mice, and in human cortical neuronal precursor cells in culture, by amyloid beta-peptide: Implications for the pathogenesis of Alzheimer's disease. *Neuromolecular Med.* 1(2): 125–135.
- Hauser, K.F., Stiene-Martin, A., Mattson, M.P., Elde, R.P., Ryan, S.E., and Godleske, C.C. (1996). mu-Opioid receptor-induced Ca²⁺ mobilization and astroglial development: Morphine inhibits DNA synthesis and stimulates cellular hypertrophy through a Ca(2+)-dependent mechanism. *Brain Res* 720(1-2): 191–203.
- Hayase, Y., Higashiyama, S., Sasahara, M., Amano, S., Nakagawa, T., Taniguchi, N. and Hazama, F. (1998). Expression of heparin-binding epidermal growth factor-like growth factor in rat brain. *Brain Res.* 784(1-2): 163–178.
- Himanen, J.P., Chumley, M.J., Lackmann, M., Li, C., Barton, W.A., Jeffrey, P.D., Vearing, C., Geleick, D., Feldheim, D.A., Boyd, A. W., Henkemeyer, M. and Nikolov, D. B. (2004). Repelling class discrimination: Ephrin-A5 binds to and activates EphB2 receptor signaling. *Nat. Neurosci.* 7(5): 501–509.
- Hitoshi, S., Alexson, T., Tropepe, V., Donoviel, D., Elia, A.J., Nye, J.S., Conlon, R.A., Mak, T.W., Bernstein, A. and van der Kooy, D. (2002). Notch pathway molecules are essential for the maintenance, but not the generation, of mammalian neural stem cells. *Genes Dev.* 16(7): 846–858.
- Hohn, A., Leibrock, J., Bailey, K. and Barde, Y.A. (1990). Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family. *Nature* 344(6264): 339–341.
- Holland, E.C., Hively, W.P., Gallo, V. and Varmus, H.E. (1998a). Modeling mutations in the G1 arrest pathway in human gliomas: Overexpression of CDK4 but not loss of INK4a-ARF induces hyperploidy in cultured mouse astrocytes. *Genes Dev.* 12(23): 3644–3649.
- Hsieh, J., Aimone, J.B., Kaspar, B.K., Kuwabara, T., Nakashima, K. and Gage, F.H. (2004). IGF-I instructs multipotent adult neural progenitor cells to become oligodendrocytes. *J. Cell. Biol.* 164(1): 111–122.
- Huang, E.J. and Reichardt, L.F. (2003). Trk receptors: Roles in neuronal signal transduction. *Annu. Rev. Biochem.* 72: 609–642.
- Ikeya, M., Lee, S.M., Johnson, J.E., McMahon, A.P. and Takada, S. (1997). Wnt signalling required for expansion of neural crest and CNS progenitors. *Nature* 389(6654): 966–970.
- Ip, N.Y. (1998). The neurotrophins and neuropoietic cytokines: Two families of growth factors acting on neural and hematopoietic cells. *Ann. N. Y. Acad. Sci* 840: 97–106.
- Ip, N.Y., Boulton, T.G., Li, Y., Verdi, J.M., Birren, S.J., Anderson, D.J. and Yancopoulos, G.D. (1994). CNTF, FGF, and NGF collaborate to drive the terminal differentiation of MAH cells into postmitotic neurons. *Neuron* 13(2): 443–455.
- Ip, N.Y., McClain, J., Barrezueta, N.X., Aldrich, T.H., Pan, L., Li, Y., Wiegand, S.J., Friedman, B., Davis, S. and Yancopoulos, G.D. (1993). The alpha component of the CNTF receptor is required for signaling and defines potential CNTF targets in the adult and during development. *Neuron* 10(1): 89–102.

- Ip, N.Y., Nye, S.H., Boulton, T.G., Davis, S., Taga, T., Li, Y., Birren, S.J., Yasukawa, K., Kishimoto, T., Anderson, D.J. and et al. (1992). CNTF and LIF act on neuronal cells via shared signaling pathways that involve the IL-6 signal transducing receptor component gp130. *Cell* 69(7): 1121–1132.
- Ip, N.Y. and Yancopoulos, G.D. (1996). The neurotrophins and CNTF: Two families of collaborative neurotrophic factors. *Annu. Rev. Neurosci.* 19: 491–515.
- Israsena, N., Hu, M., Fu, W., Kan, L. and Kessler, J.A. (2004). The presence of FGF2 signaling determines whether beta-catenin exerts effects on proliferation or neuronal differentiation of neural stem cells. *Dev. Biol.* 268(1): 220–231.
- Ivanova, N.B., Dimos, J.T., Schaniel, C., Hackney, J.A., Moore, K.A. and Lemischka, I.R. (2002). A stem cell molecular signature. *Science* 298(5593): 601–604.
- Jacks, T., Fazeli, A., Schmitt, E.M., Bronson, R.T., Goodell, M.A. and Weinberg, R.A. (1992). Effects of an Rb mutation in the mouse. *Nature* 359(6393): 295–300.
- Jacobs, J.J., Kieboom, K., Marino, S., DePinho, R.A. and van Lohuizen, M. (1999a). The oncogene and Polycomb-group gene *bmi-1* regulates cell proliferation and senescence through the *ink4a* locus. *Nature* 397(6715): 164–168.
- Jacobs, J.J., Scheijen, B., Voncken, J.W., Kieboom, K., Berns, A. and van Lohuizen, M. (1999b). *Bmi-1* collaborates with *c-Myc* in tumorigenesis by inhibiting *c-Myc*-induced apoptosis via *INK4a/ARF*. *Genes Dev.* 13(20): 2678–2690.
- Jiang, Z., Zacksenhaus, E., Gallie, B.L. and Phillips, R.A. (1997). The retinoblastoma gene family is differentially expressed during embryogenesis. *Oncogene* 14(15): 1789–1797.
- Jin, K., Mao, X.O., Sun, Y., Xie, L., Jin, L., Nishi, E., Klagsbrun, M. and Greenberg, D.A. (2002a). Heparin-binding epidermal growth factor-like growth factor: Hypoxia-inducible expression in vitro and stimulation of neurogenesis in vitro and in vivo. *J. Neurosci.* 22(13): 5365–5373.
- Jin, K., Minami, M., Lan, J.Q., Mao, X.O., Bateur, S., Simon, R.P. and Greenberg, D.A. (2001). Neurogenesis in dentate subgranular zone and rostral subventricular zone after focal cerebral ischemia in the rat. *Proc. Natl. Acad. Sci. U. S. A.* 98(8): 4710–4715.
- Jin, K., Sun, Y., Xie, L., Bateur, S., Mao, X.O., Smelick, C., Logvinova, A. and Greenberg, D.A. (2003). Neurogenesis and aging: FGF-2 and HB-EGF restore neurogenesis in hippocampus and subventricular zone of aged mice. *Aging Cell* 2(3): 175–183.
- Jin, K., Zhu, Y., Sun, Y., Mao, X.O., Xie, L. and Greenberg, D.A. (2002b). Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. *Proc. Natl. Acad. Sci. U. S. A.* 99(18): 11946–11950.
- Jin, K.L., Mao, X.O. and Greenberg, D. A. (2000a). Vascular endothelial growth factor rescues HN33 neural cells from death induced by serum withdrawal. *J. Mol. Neurosci.* 14(3): 197–203.
- Jin, K.L., Mao, X.O., Nagayama, T., Goldsmith, P.C. and Greenberg, D.A. (2000b). Induction of vascular endothelial growth factor and hypoxia-inducible factor-1 α by global ischemia in rat brain. *Neuroscience* 99(3): 577–585.
- Jori, F.P., Galderisi, U., Piegari, E., Cipollaro, M., Cascino, A., Peluso, G., Cotrufo, R., Giordano, A. and Melone, M.A. (2003). EGF-responsive rat neural stem cells: Molecular follow-up of neuron and astrocyte differentiation in vitro. *J. Cell Physiol.* 195(2): 220–233.
- Jung, A.B. and Bennett, Jr., J.P. (1996). Development of striatal dopaminergic function. I. Pre- and postnatal development of mRNAs and binding sites for

- striatal D1 (D1a) and D2 (D2a) receptors. *Brain Res. Dev. Brain Res.* 94(2): 109–120.
- Juul, S.E., Anderson, D.K., Li and Christensen, R.D. (1998). Erythropoietin and erythropoietin receptor in the developing human central nervous system. *Pediatr. Res.* 43(1): 40–49.
- Juul, S.E., Yachnis, A.T., Rojiani, A.M. and Christensen, R.D. (1999). Immunohistochemical localization of erythropoietin and its receptor in the developing human brain. *Pediatr. Dev. Pathol.* 2(2): 148–158.
- Kamijo, T., Zindy, F., Roussel, M.F., Quelle, D.E., Downing, J.R., Ashmun, R.A., Grosveld, G. and Sherr, C.J. (1997). Tumor suppression at the mouse INK4a locus mediated by the alternative reading frame product p19ARF. *Cell* 91(5): 649–659.
- Kane, C.J., Brown, G.J. and Phelan, K.D. (1996). Transforming growth factor-beta 2 both stimulates and inhibits neurogenesis of rat cerebellar granule cells in culture. *Brain Res. Dev. Brain Res.* 96(1-2): 46–51.
- Kar, S., Chabot, J.G. and Quirion, R. (1993). Quantitative autoradiographic localization of [125I]insulin-like growth factor I, [125I]insulin-like growth factor II, and [125I]insulin receptor binding sites in developing and adult rat brain. *J. Comp. Neurol.* 333(3): 375–397.
- Kastan, M.B., Zhan, Q., el-Deiry, W.S., Carrier, F., Jacks, T., Walsh, W.V., Plunkett, B.S., Vogelstein, B. and Fornace, Jr., A.J. (1992). A mammalian cell cycle checkpoint pathway utilizing p53 and GADD45 is defective in ataxia-telangiectasia. *Cell* 71(4): 587–597.
- Kawahara, N., Mishima, K., Higashiyama, S., Taniguchi, N., Tamura, A. and Kirino, T. (1999). The gene for heparin-binding epidermal growth factor-like growth factor is stress-inducible: Its role in cerebral ischemia. *J. Cereb. Blood Flow Metab.* 19(3): 307–320.
- Kelly, C.M., Zietlow, R., Dunnett, S.B. and Rosser, A.E. (2003). The effects of various concentrations of FGF-2 on the proliferation and neuronal yield of murine embryonic neural precursor cells in vitro. *Cell Transplant* 12(3): 215–223.
- Kirsch, M., Schneider, T., Lee, M.Y. and Hofmann, H.D. (1998). Lesion-induced changes in the expression of ciliary neurotrophic factor and its receptor in rat optic nerve. *Glia.* 23(3): 239–248.
- Kirschenbaum, B. and Goldman, S.A. (1995). Brain-derived neurotrophic factor promotes the survival of neurons arising from the adult rat forebrain subependymal zone. *Proc. Natl. Acad. Sci. U. S. A.* 92(1): 210–214.
- Klein, G. (1995). The extracellular matrix of the hematopoietic microenvironment. *Experientia* 51(9-10): 914–926.
- Klein, R. (2004). Eph/ephrin signaling in morphogenesis, neural development and plasticity. *Cur. Opin. Cell Biol.* 16(5): 580–589.
- Klint, P. and Claesson-Welsh, L. (1999). Signal transduction by fibroblast growth factor receptors. *Front Biosci.* 4: D165–D177.
- Koblar, S.A., Turnley, A.M., Classon, B.J., Reid, K.L., Ware, C.B., Cheema, S.S., Murphy, M. and Bartlett, P.F. (1998). Neural precursor differentiation into astrocytes requires signaling through the leukemia inhibitory factor receptor. *Proc. Natl. Acad. Sci. U. S. A.* 95(6): 3178–3181.
- Kornblum, H.I., Hussain, R.J., Bronstein, J.M., Gall, C.M., Lee, D.C. and Seroogy, K.B. (1997). Prenatal ontogeny of the epidermal growth factor receptor and its ligand, transforming growth factor alpha, in the rat brain. *J. Comp. Neurol.* 380(2): 243–261.

- Kornblum, H.I., Yanni, D.S., Easterday, M.C. and Seroogy, K.B. (2000). Expression of the EGF receptor family members ErbB2, ErbB3, and ErbB4 in germinal zones of the developing brain and in neurosphere cultures containing CNS stem cells. *Dev. Neurosci.* **22**: 16–24.
- Koury, M.J. and Bondurant, M.C. (1990a). Control of red cell production: The roles of programmed cell death (apoptosis) and erythropoietin. *Transfusion* **30**(8): 673–674.
- Koury, M.J. and Bondurant, M.C. (1990b). Erythropoietin retards DNA breakdown and prevents programmed death in erythroid progenitor cells. *Science* **248**(4953): 378–381.
- Kuhn, H.G., Winkler, J., Kempermann, G., Thal, L.J. and Gage, F.H. (1997). Epidermal growth factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain. *J. Neurosci.* **17**(15): 5820–5829.
- Kullander, K. and Klein, R. (2002). Mechanisms and functions of Eph and ephrin signalling. *Nat. Rev. Mol. Cell Biol.* **3**(7): 475–486.
- Kuppers, E. and Beyer, C., (2001). Dopamine regulates brain-derived neurotrophic factor (BDNF) expression in cultured embryonic mouse striatal cells. *Neuroreport.* **12**(6): 1175–1179.
- Kwon, C.H., Zhu, X., Zhang, J., Knoop, L.L., Tharp, R., Smeyne, R.J., Eberhart, C.G., Burger, P.C. and Baker, S.J. (2001). Pten regulates neuronal soma size: A mouse model of Lhermitte-Duclos disease. *Nat Genet.* **29**(4): 404–411.
- Lachyankar, M.B., Condon, P.J., Quesenberry, P.J., Litofsky, N.S., Recht, L.D. and Ross, A.H. (1997). Embryonic precursor cells that express Trk receptors: Induction of different cell fates by NGF, BDNF, NT-3, and CNTF. *Exp. Neurol.* **144**(2): 350–360.
- Laron, Z. (2001). Insulin-like growth factor 1 (IGF-1): A growth hormone. *Mol. Pathol.* **54**(5): 311–316.
- Lax, I., Wong, A., Lamothe, B., Lee, A., Frost, A., Hawes, J. and Schlessinger, J. (2002). The docking protein FRS2alpha controls a MAP kinase-mediated negative feedback mechanism for signaling by FGF receptors. *Mol. Cell* **10**(4): 709–719.
- Lee, E.Y., Hu, N., Yuan, S.S., Cox, L.A., Bradley, A., Lee, W.H. and Herrup, K. (1994). Dual roles of the retinoblastoma protein in cell cycle regulation and neuron differentiation. *Genes. Dev.* **8**(17): 2008–2021.
- Lee, M.H., Williams, B.O., Mulligan, G., Mukai, S., Bronson, R.T., Dyson, N., Harlow, E. and Jacks, T. (1996). Targeted disruption of p107: Functional overlap between p107 and Rb. *Genes Dev.* **10**(13): 1621–1632.
- Lee, M.Y., Deller, T., Kirsch, M., Frotscher, M. and Hofmann, H.D. (1997a). Differential regulation of ciliary neurotrophic factor (CNTF) and CNTF receptor alpha expression in astrocytes and neurons of the fascia dentata after entorhinal cortex lesion. *J. Neurosci.* **17**(3): 1137–1146.
- Lee, M.Y., Naumann, T., Kirsch, M., Frotscher, M. and Hofmann, H.D. (1997b). Transient up-regulation of ciliary neurotrophic factor receptor-alpha mRNA in axotomized rat septal neurons. *Eur. J. Neurosci.* **9**(3): 622–626.
- Lees, E., Faha, B., Dulic, V., Reed, S.I. and Harlow, E. (1992). Cyclin E/cdk2 and cyclin A/cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* **6**(10): 1874–1885.
- Leibrock, J., Lottspeich, F., Hohn, A., Hofer, M., Hengerer, B., Masiakowski, P., Thoenen H. and Barde, Y.A. (1989). Molecular cloning and expression of brain-derived neurotrophic factor. *Nature* **341**(6238): 149–152.

- Leonard, J.R., D'Sa, C., Klocke, B.J. and Roth, K.A. (2001). Neural precursor cell apoptosis and glial tumorigenesis following transplacental ethyl-nitrosourea exposure. *Oncogene* 20(57): 8281–8286.
- Leventhal, C., Rafii, S., Rafii, D., Shahar, A. and Goldman, S.A. (1999). Endothelial trophic support of neuronal production and recruitment from the adult mammalian subependyma. *Mol. Cell Neurosci.* 13(6): 450–464.
- Li, J., Yen, C., Liaw, D., Podsypanina, K., Bose, S., Wang, S.I., Puc, J., Miliaresis, C., Rodgers, L., McCombie, R., Bigner, S.H., Giovanella, B.C., Ittmann, M., Tycko, B., Hibshoosh, H., Wigler, M. H. and Parsons, R.,(1997). PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 275(5308): 1943–1947.
- Li, L., Liu, F. and Ross, A.H. (2003). PTEN regulation of neural development and CNS stem cells. *J. Cell Biochem.* 88(1): 24–28.
- Li, M., Sendtner, M. and Smith, A. (1995). Essential function of LIF receptor in motor neurons. *Nature* 378(6558): 724–727.
- Li, W., Cogswell, C.A. and LoTurco, J.J. (1998). Neuronal differentiation of precursors in the neocortical ventricular zone is triggered by BMP. *J. Neurosci.* 18(21): 8853–8862.
- Li, Y., Graham, C., Lacy, S., Duncan, A.M. and Whyte, P. (1993). The adenovirus E1A-associated 130-kD protein is encoded by a member of the retinoblastoma gene family and physically interacts with cyclins A and E. *Genes Dev.* 7(12A): 2366–2377.
- Liem, K.F., Jr., Tremml, G. and Jessell, T.M. (1997). A role for the roof plate and its resident TGFbeta-related proteins in neuronal patterning in the dorsal spinal cord. *Cell* 91(1): 127–138.
- Liem, K.F., Jr., Tremml, G., Roelink, H. and Jessell, T.M. (1995). Dorsal differentiation of neural plate cells induced by BMP-mediated signals from epidermal ectoderm. *Cell* 82(6): 969–979.
- Lillien, L. and Raphael, H. (2000). BMP and FGF regulate the development of EGF-responsive neural progenitor cells. *Development* 127(22): 4993–5005.
- Lim, D.A., Tramontin, A.D., Trevejo, J.M., Herrera, D.G., Garcia-Verdugo, J.M. and Alvarez-Buylla, A. (2000). Noggin antagonizes BMP signaling to create a niche for adult neurogenesis. *Neuron* 28(3): 713–726.
- Lin, C.S., Lim, S.K., D'Agati, V. and Costantini, F. (1996). Differential effects of an erythropoietin receptor gene disruption on primitive and definitive erythropoiesis. *Genes Dev.* 10(2): 154–164.
- Lindsell, C.E., Boulter, J., diSibio, G., Gossler, A. and Weinmaster, G. (1996). Expression patterns of Jagged, Delta1, Notch1, Notch2, and Notch3 genes identify ligand-receptor pairs that may function in neural development. *Mol. Cell Neurosci.* 8(1): 14–27.
- Liu, J., Solway, K., Messing, R.O. and Sharp, F.R. (1998). Increased neurogenesis in the dentate gyrus after transient global ischemia in gerbils. *J. Neurosci.* 18(19): 7768–7778.
- Liu, J.P., Baker, J., Perkins, A.S., Robertson, E.J. and Efstratiadis, A. (1993). Mice carrying null mutations of the genes encoding insulin-like growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). *Cell* 75(1): 59–72.
- Liu, Z.Y., Chin, K. and Noguchi, C.T. (1994). Tissue specific expression of human erythropoietin receptor in transgenic mice. *Dev. Biol.* 166(1): 159–169.
- Logan, C.Y. and Nusse, R. (2004). The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev. Biol.* 20: 781–810.

- Luo, R.X., Postigo, A.A. and Dean, D.C. (1998). Rb interacts with histone deacetylase to repress transcription. *Cell* 92(4): 463–473.
- Luskin, M.B. and Coskun, V. (2002). The progenitor cells of the embryonic telencephalon and the neonatal anterior subventricular zone differentially regulate their cell cycle. *Chem. Senses* 27(6): 577–580.
- Machold, R., Hayashi, S., Rutlin, M., Muzumdar, M.D., Nery, S., Corbin, J.G., Gritli-Linde, A., Dellovade, T., Porter, J.A., Rubin, L.L., Dudek, H., McMahon, A.P. and Fishell, G. (2003). Sonic hedgehog is required for progenitor cell maintenance in telencephalic stem cell niches. *Neuron* 39(6): 937–950.
- MacLennan, A.J., Vinson, E.N., Marks, L., McLaurin, D.L., Pfeifer, M. and Lee, N. (1996). Immunohistochemical localization of ciliary neurotrophic factor receptor alpha expression in the rat nervous system. *J. Neurosci.* 16(2): 621–630.
- Magavi, S.S., Leavitt, B.R. and Macklis, J.D. (2000). Induction of neurogenesis in the neocortex of adult mice. *Nature* 405(6789): 951–955.
- Magnaghi-Jaulin, L., Groisman, R., Naguibneva, I., Robin, P., Lorain, S., Le Villain, J.P., Troalen, F., Trouche, D. and Harel-Bellan, A. (1998). Retinoblastoma protein represses transcription by recruiting a histone deacetylase. *Nature* 391(6667): 601–65.
- Mahanthappa, N.K. and Schwarting, G.A. (1993). Peptide growth factor control of olfactory neurogenesis and neuron survival in vitro: Roles of EGF and TGF-beta s. *Neuron* 10(2): 293–305.
- Maisonpierre, P.C., Belluscio, L., Squinto, S., Ip, N.Y., Furth, M.E., Lindsay, R.M. and Yancopoulos, G.D. (1990). Neurotrophin-3: A neurotrophic factor related to NGF and BDNF. *Science* 247(4949 Pt 1): 1446–1451.
- Malberg, J.E., Eisch, A.J., Nestler, E.J. and Duman, R.S. (2000). Chronic antidepressant treatment increases neurogenesis in adult rat hippocampus. *J. Neurosci.* 20(24): 9104–9110.
- Mallamaci, A., Iannone, R., Briata, P., Pintonello, L., Mercurio, S., Boncinelli, E. and Corte, G. (1998). EMX2 protein in the developing mouse brain and olfactory area. *Mech. Dev.* 77(2): 165–172.
- Mallamaci, A., Mercurio, S., Muzio, L., Cecchi, C., Pardini, C.L., Gruss, P. and Boncinelli, E. (2000). The lack of Emx2 causes impairment of Reelin signaling and defects of neuronal migration in the developing cerebral cortex. *J. Neurosci.* 20(3): 1109–1118.
- Marmur, R., Kessler, J.A., Zhu, G., Gokhan, S. and Mehler, M.F. (1998). Differentiation of oligodendroglial progenitors derived from cortical multipotent cells requires extrinsic signals including activation of gp130/LIFbeta receptors. *J. Neurosci.* 18(23): 9800–9811.
- Martens, D.J., Tropepe, V. and van Der Kooy, D. (2000). Separate proliferation kinetics of fibroblast growth factor-responsive and epidermal growth factor-responsive neural stem cells within the embryonic forebrain germinal zone. *J. Neurosci.* 20(3): 1085–1095.
- Marti, E., Bumcrot, D.A., Takada, R. and McMahon, A.P. (1995). Requirement of 19K form of Sonic hedgehog for induction of distinct ventral cell types in CNS explants. *Nature* 375(6529): 322–325.
- Mason, I., Chambers, D., Shamim, H., Walshe, J. and Irving, C. (2000). Regulation and function of FGF8 in patterning of midbrain and anterior hindbrain. *Biochem. Cell Biol.* 78(5): 577–584.

- Masu, Y., Wolf, E., Holtmann, B., Sendtner, M., Brem, G. and Thoenen, H. (1993). Disruption of the CNTF gene results in motor neuron degeneration. *Nature* 365(6441): 27–32.
- Matsuoka, S., Edwards, M.C., Bai, C., Parker, S., Zhang, P., Baldini, A., Harper, J.W. and Elledge, S.J. (1995). p57KIP2, a structurally distinct member of the p21CIP1 Cdk inhibitor family, is a candidate tumor suppressor gene. *Genes Dev.* 9(6): 650–662.
- Matsuzaki, H., Tamatani, M., Yamaguchi, A., Namikawa, K., Kiyama, H., Vitek, M.P., Mitsuda, N. and Tohyama, M. (2001). Vascular endothelial growth factor rescues hippocampal neurons from glutamate-induced toxicity: Signal transduction cascades. *Faseb. J.* 15(7): 1218–1220.
- Mattson, M.P., Maudsley, S. and Martin, B. (2004). BDNF and 5-HT: A dynamic duo in age-related neuronal plasticity and neurodegenerative disorders. *Trends Neurosci.* 27(10): 589–594.
- McEwen, B., Akama, K., Alves, S., Brake, W.G., Bulloch, K., Lee, S., Li, C., Yuen, G. and Milner, T. A. (2001). Tracking the estrogen receptor in neurons: Implications for estrogen-induced synapse formation. *Proc. Natl. Acad. Sci. U. S. A.* 98(13): 7093–7100.
- Megason, S.G. and McMahon, A.P. (2002). A mitogen gradient of dorsal midline Wnts organizes growth in the CNS. *Development* 129(9): 2087–2098.
- Menezes, J.R., Dias, F., Garson, A.V. and Lent, R. (1998). Restricted distribution of S-phase cells in the anterior subventricular zone of the postnatal mouse forebrain. *Anat. Embryol. (Berl.)* 198(3): 205–211.
- Menezes, J.R., Smith, C.M., Nelson, K.C. and Luskin, M.B. (1995). The division of neuronal progenitor cells during migration in the neonatal mammalian forebrain. *Mol. Cell Neurosci.* 6(6): 496–508.
- Mercier, F., Kitasako, J.T. and Hatton, G.I. (2002). Anatomy of the brain neurogenic zones revisited: Fractones and the fibroblast/macrophage network. *J. Comp. Neurol.* 451(2): 170–188.
- Millan, F.A., Denhez, F., Kondaiiah, P. and Akhurst, R.J. (1991). Embryonic gene expression patterns of TGF beta 1, beta 2 and beta 3 suggest different developmental functions in vivo. *Development* 111(1): 131–143.
- Mishima, K., Higashiyama, S., Nagashima, Y., Miyagi, Y., Tamura, A., Kawahara, N., Taniguchi, N., Asai, A., Kuchino, Y. and Kirino, T. (1996). Regional distribution of heparin-binding epidermal growth factor-like growth factor mRNA and protein in adult rat forebrain. *Neurosci. Lett.* 213(3): 153–156.
- Moghaddam, B., Bolinao, M.L., Stein-Behrens, B. and Sapolsky, R. (1994). Glucocorticoids mediate the stress-induced extracellular accumulation of glutamate. *Brain Res.* 655(1-2): 251–254.
- Molofsky, A.V., Pardal, R., Iwashita, T., Park, I.K., Clarke, M.F. and Morrison, S.J. (2003). Bmi-1 dependence distinguishes neural stem cell self-renewal from progenitor proliferation. *Nature* 425(6961): 962–967.
- Morrison, S.J., Csete, M., Groves, A.K., Melega, W., Wold, B. and Anderson, D.J. (2000a). Culture in reduced levels of oxygen promotes clonogenic sympathoadrenal differentiation by isolated neural crest stem cells. *J. Neurosci.* 20(19): 7370–7376.
- Morrison, S.J., Perez, S.E., Qiao, Z., Verdi, J.M., Hicks, C., Weinmaster, G. and Anderson, D.J. (2000b). Transient Notch activation initiates an irreversible switch from neurogenesis to gliogenesis by neural crest stem cells. *Cell* 101(5): 499–510.

- Morrow, T., Song, M.R. and Ghosh, A. (2001). Sequential specification of neurons and glia by developmentally regulated extracellular factors. *Development* 128(18): 3585–3594.
- Morshead, C.M., Craig, C.G. and van der Kooy, D. (1998). In vivo clonal analyses reveal the properties of endogenous neural stem cell proliferation in the adult mammalian forebrain. *Development* 125(12): 2251–2261.
- Morshead, C.M., Reynolds, B.A., Craig, C.G., McBurney, M.W., Staines, W.A., Morassutti, D., Weiss, S. and van der Kooy, D. (1994). Neural stem cells in the adult mammalian forebrain: A relatively quiescent subpopulation of subependymal cells. *Neuron* 13(5): 1071–1082.
- Morshead, C.M. and van der Kooy, D. (1992). Postmitotic death is the fate of constitutively proliferating cells in the subependymal layer of the adult mouse brain. *J. Neurosci.* 12(1): 249–256.
- Mundle, S.D. and Saberwal, G. (2003). Evolving intricacies and implications of E2F1 regulation. *Faseb. J.* 17(6): 569–574.
- Nakagawa, T., Sasahara, M., Hayase, Y., Haneda, M., Yasuda, H., Kikkawa, R., Higashiyama, S. and Hazama, F. (1998). Neuronal and glial expression of heparin-binding EGF-like growth factor in central nervous system of prenatal and early-postnatal rat. *Brain Res. Dev. Brain Res.* 108(1-2): 263–262.
- Nakamura, Y., Sakakibara, S., Miyata, T., Ogawa, M., Shimazaki, T., Weiss, S., Kageyama, R. and Okano, H. (2000). The bHLH gene *hes1* as a repressor of the neuronal commitment of CNS stem cells. *J. Neurosci.* 20(1): 283–293.
- Nakashima, K., Wiese, S., Yanagisawa, M., Arakawa, H., Kimura, N., Hisatsune, T., Yoshida, K., Kishimoto, T., Sendtner, M. and Taga, T. (1999a). Developmental requirement of gp130 signaling in neuronal survival and astrocyte differentiation. *J. Neurosci.* 19(13): 5429–5434.
- Nakashima, K., Yanagisawa, M., Arakawa, H., Kimura, N., Hisatsune, T., Kawabata, M., Miyazono, K. and Taga, T. (1999b). Synergistic signaling in fetal brain by STAT3-Smad1 complex bridged by p300. *Science* 284(5413): 479–482.
- Nandurkar, H.H., Hilton, D.J., Nathan, P., Willson, T., Nicola, N. and Begley, C.G. (1996). The human IL-11 receptor requires gp130 for signalling: Demonstration by molecular cloning of the receptor. *Oncogene* 12(3): 585–593.
- Nevins, J.R. (2001). The Rb/E2F pathway and cancer. *Hum Mol Genet* 10(7): 699–703.
- Nieouillon, A. (2002). Dopamine and the regulation of cognition and attention. *Prog Neurobiol* 67(1): 53–83.
- Nowakowski, R.S., Caviness, Jr., V.S., Takahashi, T. and Hayes, N.L. (2002). Population dynamics during cell proliferation and neuronogenesis in the developing murine neocortex. *Results Probl Cell Differ* 39: 1–25.
- O’Leary, D.D. and Wilkinson, D.G. (1999). Eph receptors and ephrins in neural development. *Curr. Opin. Neurobiol.* 9(1): 65–73.
- Ohmiya, M., Fukumitsu, H., Nitta, A., Nomoto, H., Furukawa, Y. and Furukawa, S. (2001). Administration of FGF-2 to embryonic mouse brain induces hydrocephalic brain morphology and aberrant differentiation of neurons in the postnatal cerebral cortex. *J. Neurosci. Res.* 65(3): 228–235.
- Ohsawa, I., Takamura, C., Morimoto, T., Ishiguro, M. and Kohsaka, S. (1999). Amino-terminal region of secreted form of amyloid precursor protein stimulates proliferation of neural stem cells. *Eur. J. Neurosci.* 11(6): 1907–1913.

- Ohtani, N., Goto, T., Waeber, C. and Bhide, P.G. (2003). Dopamine modulates cell cycle in the lateral ganglionic eminence. *J. Neurosci.* 23(7): 2840–2850.
- Opanashuk, L.A. and Hauser, K.F. (1998). Opposing actions of the EGF family and opioids: Heparin binding-epidermal growth factor (HB-EGF) protects mouse cerebellar neuroblasts against the antiproliferative effect of morphine. *Brain Res.* 804(1): 87–94.
- Orioli, D. and Klein, R. (1997). The Eph receptor family: Axonal guidance by contact repulsion. *Trends Genet.* 13(9): 354–359.
- Orr-Urtreger, A., Givol, D., Yayon, A., Yarden, Y. and Lonai, P. (1991). Developmental expression of two murine fibroblast growth factor receptors, flg and bek. *Development* 113(4): 1419–1434.
- Ortega, S., Malumbres, M. and Barbacid, M. (2002). Cyclin D-dependent kinases, INK4 inhibitors and cancer. *Biochim. Biophys. Acta.* 1602(1): 73–87.
- Otero, J.J., Fu, W., Kan, L., Cuadra, A.E. and Kessler, J.A. (2004). Beta-catenin signaling is required for neural differentiation of embryonic stem cells. *Development* 131(15): 3545–3557.
- Paggi, M.G., Baldi, A., Bonetto, F. and Giordano, A. (1996). Retinoblastoma protein family in cell cycle and cancer: A review. *J. Cell. Biochem.* 62(3): 418–430.
- Palma, V., Lim, D.A., Dahmane, N., Sanchez, P., Brionne, T.C., Herzberg, C.D., Gitton, Y., Carleton, A., Alvarez-Buylla, A. and Altaba, A.R. (2005). Sonic hedgehog controls stem cell behavior in the postnatal and adult brain. *Development* 132(2): 335–344.
- Palmer, A. and Klein, R. (2003). Multiple roles of ephrins in morphogenesis, neuronal networking, and brain function. *Genes Dev.* 17(12): 1429–1450.
- Palmert, M.R., Podlisny, M.B., Witker, D.S., Oltersdorf, T., Younkin, L.H., Selkoe, D.J. and Younkin, S.G. (1989). The beta-amyloid protein precursor of Alzheimer disease has soluble derivatives found in human brain and cerebrospinal fluid. *Proc. Natl. Acad. Sci. U. S. A.* 86(16): 6338–6342.
- Panchision, D.M. and McKay, R.D. (2002). The control of neural stem cells by morphogenic signals. *Curr. Opin. Genet. Dev.* 12(4): 478–487.
- Panchision, D.M., Pickel, J.M., Studer, L., Lee, S.H., Turner, P.A., Hazel, T.G. and McKay, R.D. (2001). Sequential actions of BMP receptors control neural precursor cell production and fate. *Genes Dev.* 15(16): 2094–2110.
- Parent, J.M., Yu, T.W., Leibowitz, R.T., Geschwind, D.H., Sloviter, R.S. and Lowenstein, D.H. (1997). Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J. Neurosci.* 17(10): 3727–3738.
- Park, J.K., Williams, B.P., Alberta, J.A. and Stiles, C.D. (1999). Bipotent cortical progenitor cells process conflicting cues for neurons and glia in a hierarchical manner. *J. Neurosci.* 19(23): 10383–10389.
- Partanen, J., Schwartz, L. and Rossant, J. (1998). Opposite phenotypes of hypomorphic and Y766 phosphorylation site mutations reveal a function for Fgfr1 in anteroposterior patterning of mouse embryos. *Genes Dev.* 12(15): 2332–2344.
- Pasquale, E.B. (2004). Eph-ephrin promiscuity is now crystal clear. *Nat. Neurosci.* 7(5): 417–418.
- Pellegrini, M., Mansouri, A., Simeone, A., Boncinelli, E. and Gruss, P. (1996). Dentate gyrus formation requires Emx2. *Development* 122(12): 3893–3898.
- Pencea, V., Bingaman, K.D., Wiegand, S.J. and Luskin, M.B. (2001). Infusion of brain-derived neurotrophic factor into the lateral ventricle of the adult rat leads to

- new neurons in the parenchyma of the striatum, septum, thalamus, and hypothalamus. *J. Neurosci.* 21(17): 6706–6717.
- Pennica, D., Shaw, K.J., Swanson, T.A., Moore, M.W., Shelton, D.L., Zioncheck, K.A., Rosenthal, A., Taga, T., Paoni, N.F. and Wood, W.I. (1995). Cardiotropin-1. Biological activities and binding to the leukemia inhibitory factor receptor/gp130 signaling complex. *J. Biol. Chem.* 270(18): 10915–22.
- Pesce, M., Farrace, M.G., Piacentini, M., Dolci, S. and De Felici, M. (1993). Stem cell factor and leukemia inhibitory factor promote primordial germ cell survival by suppressing programmed cell death (apoptosis). *Development* 118(4): 1089–1094.
- Pilon, C., Levesque, D., Dimitriadou, V., Griffon, N., Martres, M.P., Schwartz, J.C. and Sokoloff, P. 1994. Functional coupling of the human dopamine D3 receptor in a transfected NG 108-15 neuroblastoma-glioma hybrid cell line. *Eur. J. Pharmacol.* 268(2): 129–139.
- Polyak, K., Lee, M.H., Erdjument-Bromage, H., Koff, A., Roberts, J.M., Tempst, P. and Massague, J. (1994). Cloning of p27Kip1, a cyclin-dependent kinase inhibitor and a potential mediator of extracellular antimitogenic signals. *Cell* 78(1): 59–66.
- Puymirat, J., Mieke, M., Marchand, R., Sarlieve, L. and Dussault, J.H. (1991). Immunocytochemical localization of thyroid hormone receptors in the adult rat brain. *Thyroid* 1(2): 173–184.
- Qian, X., Davis, A.A., Goderie, S.K. and Temple, S. (1997). FGF2 concentration regulates the generation of neurons and glia from multipotent cortical stem cells. *Neuron* 18(1): 81–93.
- Qian, X., Goderie, S.K., Shen, Q., Stern, J.H. and Temple, S. (1998). Intrinsic programs of patterned cell lineages in isolated vertebrate CNS ventricular zone cells. *Development* 125(16): 3143–3152.
- Qian, X., Shen, Q., Goderie, S.K., He, W., Capela, A., Davis, A.A. and Temple, S. (2000). Timing of CNS cell generation: A programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. *Neuron* 28(1): 69–80.
- Qiu, J., Takagi, Y., Harada, J., Rodrigues, N., Moskowitz, M.A., Scadden, D.T. and Cheng, T. (2004). Regenerative response in ischemic brain restricted by p21cip1/waf1. *J. Exp. Med.* 199(7): 937–945.
- Quelle, D.E., Cheng, M., Ashmun, R.A. and Sherr, C.J. (1997). Cancer-associated mutations at the INK4a locus cancel cell cycle arrest by p16INK4a but not by the alternative reading frame protein p19ARF. *Proc. Natl. Acad. Sci. U. S. A.* 94(2): 669–673.
- Quelle, D.E., Zindy, F., Ashmun, R.A. and Sherr, C.J. (1995). Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest. *Cell* 83(6): 993–1000.
- Raballo, R., Rhee, J., Lyn-Cook, R., Leckman, J.F., Schwartz, M.L. and Vaccarino, F.M. (2000). Basic fibroblast growth factor (Fgf2) is necessary for cell proliferation and neurogenesis in the developing cerebral cortex. *J. Neurosci.* 20(13): 5012–5023.
- Rajan, P. and McKay, R.D. (1998). Multiple routes to astrocytic differentiation in the CNS. *J. Neurosci.* 18(10): 3620–3629.
- Rajan, P., Panchision, D.M., Newell, L.F. and McKay, R.D. (2003). BMPs signal alternately through a SMAD or FRAP-STAT pathway to regulate fate choice in CNS stem cells. *J. Cell Biol.* 161(5): 911–921.
- Recht, L., Jang, T., Savarese, T. and Litofsky, N. S. (2003). Neural stem cells and neuro-oncology: Quo vadis? *J. Cell Biochem.* 88(1): 11–19.

- Reissmann, E., Ernsberger, U., Francis-West, P. H., Rueger, D., Brickell, P.M. and Rohrer, H. (1996). Involvement of bone morphogenetic protein-4 and bone morphogenetic protein-7 in the differentiation of the adrenergic phenotype in developing sympathetic neurons. *Development* 122(7): 2079–2088.
- Represa, J., Leon, Y., Miner, C. and Giraldez, F. (1991). The int-2 proto-oncogene is responsible for induction of the inner ear. *Nature* 353(6344): 561–563.
- Reuss, B. and von Bohlen und Halbach, O. (2003). Fibroblast growth factors and their receptors in the central nervous system. *Cell Tissue Res.* 313(2): 139–157.
- Reynolds, B.A. and Weiss, S. (1992). Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255(5052): 1707–1710.
- Risau, W. (1997). Mechanisms of angiogenesis. *Nature* 386(6626): 671–674.
- Roelink, H., Porter, J.A., Chiang, C., Tanabe, Y., Chang, D.T., Beachy, P.A. and Jessell, T.M. (1995). Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell* 81(3): 445–455.
- Roof, R.L. and Hall, E.D. (2000). Gender differences in acute CNS trauma and stroke: Neuroprotective effects of estrogen and progesterone. *J. Neurotrauma* 17(5): 367–388.
- Rudge, J.S., Eaton, M.J., Mather, P., Lindsay, R.M. and Whittemore, S.R. (1996). CNTF induces raphe neuronal precursors to switch from a serotonergic to a cholinergic phenotype in vitro. *Mol. Cell Neurosci.* 7(3): 204–221.
- Russo, A.A., Jeffrey, P.D., Patten, A.K., Massague, J. and Pavletich, N.P. (1996). Crystal structure of the p27Kip1 cyclin-dependent-kinase inhibitor bound to the cyclin A-Cdk2 complex. *Nature* 382(6589): 325–331.
- Saadat, S., Sendtner, M. and Rohrer, H. (1989). Ciliary neurotrophic factor induces cholinergic differentiation of rat sympathetic neurons in culture. *J. Cell. Biol.* 108(5): 1807–1816.
- Santarelli, L., Saxe, M., Gross, C., Surget, A., Battaglia, F., Dulawa, S., Weisstaub, N., Lee, J., Duman, R., Arancio, O., Belzung, C. and Hen, R. (2003). Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 301(5634): 805–809.
- Schmid, P., Cox, D., Bilbe, G., Maier, R. and McMaster, G.K. (1991). Differential expression of TGF beta 1, beta 2 and beta 3 genes during mouse embryogenesis. *Development* 111(1): 117–130.
- Schultze, B. and Korr, H., (1981). Cell kinetic studies of different cell types in the developing and adult brain of the rat and the mouse: A review. *Cell Tissue Kinet* 14(3): 309–325.
- See, J., Zhang, X., Eraydin, N., Mun, S.B., Mamontov, P., Golden, J.A. and Grinspan, J.B. (2004). Oligodendrocyte maturation is inhibited by bone morphogenetic protein. *Mol. Cell Neurosci.* 26(4): 481–492.
- Sendtner, M., Dittrich, F., Hughes, R.A. and Thoenen, H. (1994). Actions of CNTF and neurotrophins on degenerating motoneurons: Preclinical studies and clinical implications. *J. Neuro. Sci.* 124 Suppl: 77–83.
- Seniuk-Tatton, N.A., Henderson, J.T. and Roder, J.C. (1995). Neurons express ciliary neurotrophic factor mRNA in the early postnatal and adult rat brain. *J. Neurosci. Res.* 41(5): 663–676.

- Shah, N.M., Groves, A.K. and Anderson, D.J. (1996). Alternative neural crest cell fates are instructively promoted by TGFbeta superfamily members. *Cell* 85(3): 331–343.
- Shen, Q., Goderie, S.K., Jin, L., Karanth, N., Sun, Y., Abramova, N., Vincent, P., Pumiglia, K. and Temple, S. (2004). Endothelial cells stimulate self-renewal and expand neurogenesis of neural stem cells. *Science* 304(5675): 1338–1340.
- Sherr, C.J. (1994). G1 phase progression: Cycling on cue. *Cell* 79(4): 551–555.
- Sherr, C.J. and Roberts, J.M. (1995). Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev.* 9(10): 1149–1163.
- Sherr, C.J. and Roberts, J.M. (1999). CDK inhibitors: Positive and negative regulators of G1-phase progression. *Genes Dev.* 13(12): 1501–1512.
- Shi, Y. and Massague, J. (2003). Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 113(6): 685–700.
- Shimazaki, T., Shingo, T. and Weiss, S. (2001). The ciliary neurotrophic factor/leukemia inhibitory factor/gp130 receptor complex operates in the maintenance of mammalian forebrain neural stem cells. *J. Neurosci.* 21(19): 7642–7653.
- Shingo, T., Gregg, C., Enwere, E., Fujikawa, H., Hassam, R., Geary, C., Cross, J.C. and Weiss, S. (2003). Pregnancy-stimulated neurogenesis in the adult female forebrain mediated by prolactin. *Science* 299(5603): 117–120.
- Shingo, T., Sorokan, S.T., Shimazaki, T. and Weiss, S. (2001). Erythropoietin regulates the in vitro and in vivo production of neuronal progenitors by mammalian forebrain neural stem cells. *J. Neurosci.* 21(24): 9733–9743.
- Shughrue, P.J., Stumpf, W.E., MacLusky, N.J., Zielinski, J.E. and Hochberg, R.B. (1990). Developmental changes in estrogen receptors in mouse cerebral cortex between birth and postweaning: Studied by autoradiography with 11 beta-methoxy-16 alpha-[125I]iodoestradiol. *Endocrinology* 126(2): 1112–1124.
- Sibilia, M., Steinbach, J.P., Stingl, L., Aguzzi, A. and Wagner, E.F. (1998). A strain-independent postnatal neurodegeneration in mice lacking the EGF receptor. *Embo. J* 17(3): 719–731.
- Sibilia, M. and Wagner, E.F. (1995). Strain-dependent epithelial defects in mice lacking the EGF receptor. *Science* 269(5221): 234–238.
- Silverman, W.F., Krum, J.M., Mani, N. and Rosenstein, J.M. (1999). Vascular, glial and neuronal effects of vascular endothelial growth factor in mesencephalic explant cultures. *Neuroscience* 90(4): 1529–1541.
- Sleeman, M.W., Anderson, K.D., Lambert, P.D., Yancopoulos, G.D. and Wiegand, S.J. (2000). The ciliary neurotrophic factor and its receptor, CNTFR alpha. *Pharm. Acta. Helv.* 74(2-3): 265–272.
- Smith, C.M. and Luskin, M.B. (1998). Cell cycle length of olfactory bulb neuronal progenitors in the rostral migratory stream. *Dev. Dyn.* 213(2): 220–227.
- Sokoloff, P., Guillin, O., Diaz, J., Carroll, P. and Griffon, N. (2002). Brain-derived neurotrophic factor controls dopamine D3 receptor expression: Implications for neurodevelopmental psychiatric disorders. *Neurotox Res.* 4(7-8): 671–678.
- Sondell, M., Lundborg, G. and Kanje, M. (1999). Vascular endothelial growth factor has neurotrophic activity and stimulates axonal outgrowth, enhancing cell survival and Schwann cell proliferation in the peripheral nervous system. *J. Neurosci.* 19(14): 5731–5740.
- Sondell, M., Sundler, F. and Kanje, M., (2000). Vascular endothelial growth factor is a neurotrophic factor which stimulates axonal outgrowth through the flk-1 receptor. *Eur. J. Neurosci.* 12(12): 4243–4254.

- Song, M.R. and Ghosh, A. (2004). FGF2-induced chromatin remodeling regulates CNTF-mediated gene expression and astrocyte differentiation. *Nat. Neurosci.* 7(3): 229–235.
- Soria, J.M., Tagliatela, P., Gil-Perotin, S., Galli, R., Gritti, A., Verdugo, J.M. and Bertuzzi, S. (2004). Defective postnatal neurogenesis and disorganization of the rostral migratory stream in absence of the *Vax1* homeobox gene. *J. Neurosci.* 24(49): 11171–11181.
- Stein-Behrens, B.A., Lin, W.J. and Sapolsky, R.M. (1994). Physiological elevations of glucocorticoids potentiate glutamate accumulation in the hippocampus. *J. Neurochem.* 63(2): 596–602.
- Stiene-Martin, A., Knapp, P.E., Martin, K., Gurwell, J.A., Ryan, S., Thornton, S.R., Smith, F.L. and Hauser, K.F. (2001). Opioid system diversity in developing neurons, astroglia, and oligodendroglia in the subventricular zone and striatum: Impact on gliogenesis in vivo. *Glia.* 36(1): 78–88.
- Struder, H.K. and Weicker, H. (2001a). Physiology and pathophysiology of the serotonergic system and its implications on mental and physical performance. Part I. *Int. J. Sports Med.* 22(7): 467–481.
- Struder, H.K. and Weicker, H. (2001b). Physiology and pathophysiology of the serotonergic system and its implications on mental and physical performance. Part II. *Int. J. Sports Med.* 22(7): 482–497.
- Stuckmann, I., Weigmann, A., Shevchenko, A., Mann, M. and Huttner, W.B. (2001). Ephrin B1 is expressed on neuroepithelial cells in correlation with neocortical neurogenesis. *J. Neurosci.* 21(8): 2726–2737.
- Studer, L., Csete, M., Lee, S.H., Kabbani, N., Walikonis, J., Wold, B. and McKay, R. (2000). Enhanced proliferation, survival, and dopaminergic differentiation of CNS precursors in lowered oxygen. *J. Neurosci.* 20(19): 7377–7383.
- Sun, Y., Nadal-Vicens, M., Misono, S., Lin, M.Z., Zubiaga, A., Hua, X., Fan, G. and Greenberg, M.E. (2001). Neurogenin promotes neurogenesis and inhibits glial differentiation by independent mechanisms. *Cell* 104(3): 365–376.
- Takagi, Y., Nozaki, K., Takahashi, J., Yodoi, J., Ishikawa, M. and Hashimoto, N. (1999). Proliferation of neuronal precursor cells in the dentate gyrus is accelerated after transient forebrain ischemia in mice. *Brain Res.* 831(1-2): 283–287.
- Takahashi, T., Nowakowski, R.S. and Caviness, Jr., V.S. (1996). The leaving or Q fraction of the murine cerebral proliferative epithelium: A general model of neocortical neurogenesis. *J. Neurosci.* 16(19): 6183–6196.
- Tamura, M., Gu, J., Matsumoto, K., Aota, S., Parsons, R. and Yamada, K.M. (1998). Inhibition of cell migration, spreading, and focal adhesions by tumor suppressor PTEN. *Science* 280(5369): 1614–1617.
- Tanapat, P., Hastings, N.B. and Gould, E. (2005). Ovarian steroids influence cell proliferation in the dentate gyrus of the adult female rat in a dose- and time-dependent manner. *J. Comp. Neurol.* 481(3): 252–265.
- Tanapat, P., Hastings, N.B., Reeves, A.J. and Gould, E. (1999). Estrogen stimulates a transient increase in the number of new neurons in the dentate gyrus of the adult female rat. *J. Neurosci.* 19(14): 5792–5801.
- Tao, Y., Black, I.B. and DiCicco-Bloom, E. (1996). Neurogenesis in neonatal rat brain is regulated by peripheral injection of basic fibroblast growth factor (bFGF). *J. Comp. Neurol.* 376(4): 653–663.
- Temple, S. (2001). Stem cell plasticity—building the brain of our dreams. *Nat. Rev. Neurosci.* 2(7): 513520.

- Teng, K.K. and Hempstead, B.L. (2004). Neurotrophins and their receptors: Signaling trios in complex biological systems. *Cell Mol. Life Sci.* 61(1): 35–48.
- Threadgill, D.W., Dlugosz, A.A., Hansen, L.A., Tennenbaum, T., Lichti, U., Yee, D., LaMantia, C., Mourton, T., Herrup, K., Harris, R.C. and et al. (1995). Targeted disruption of mouse EGF receptor: Effect of genetic background on mutant phenotype. *Science* 269(5221): 230–234.
- Tole, S., Goudreau, G., Assimacopoulos, S. and Grove, E.A. (2000). *Emx2* is required for growth of the hippocampus but not for hippocampal field specification. *J. Neurosci.* 20(7): 2618–2625.
- Toyoshima, H. and Hunter, T. (1994). p27, a novel inhibitor of G1 cyclin-Cdk protein kinase activity, is related to p21. *Cell* 78(1): 67–74.
- Tropepe, V., Craig, C.G., Morshead, C.M. and van der Kooy, D. (1997). Transforming growth factor- α null and senescent mice show decreased neural progenitor cell proliferation in the forebrain subependyma. *J. Neurosci.* 17(20): 7850–7859.
- Tropepe, V., Sibilian, M., Ciruna, B.G., Rossant, J., Wagner, E.F. and van der Kooy, D. (1999). Distinct neural stem cells proliferate in response to EGF and FGF in the developing mouse telencephalon. *Dev. Biol.* 208(1): 166–188.
- Turnley, A.M. and Bartlett, P.F. (2000). Cytokines that signal through the leukemia inhibitory factor receptor- β complex in the nervous system. *J. Neurochem.* 74(3): 889–899.
- Uhrbom, L., Dai, C., Celestino, J.C., Rosenblum, M.K., Fuller, G.N. and Holland, E.C. (2002). *Ink4a-Arf* loss cooperates with *KRas* activation in astrocytes and neural progenitors to generate glioblastomas of various morphologies depending on activated Akt. *Cancer Res.* 62(19): 5551–5558.
- Vaccarino, F.M., Schwartz, M.L., Hartigan, D. and Leckman, J.F. (1995). Basic fibroblast growth factor increases the number of excitatory neurons containing glutamate in the cerebral cortex. *Cereb Cortex* 5(1): 64–78.
- Vaccarino, F.M., Schwartz, M.L., Raballo, R., Nilsen, J., Rhee, J., Zhou, M., Doetschman, T., Coffin, J.D., Wyland, J.J. and Hung, Y.T. (1999a). Changes in cerebral cortex size are governed by fibroblast growth factor during embryogenesis. *Nat. Neurosci.* 2(3): 246–253.
- Vaccarino, F.M., Schwartz, M.L., Raballo, R., Rhee, J. and Lyn-Cook, R. (1999b). Fibroblast growth factor signaling regulates growth and morphogenesis at multiple steps during brain development. *Curr. Top Dev. Biol.* 46: 179–200.
- Van Kampen, J.M., Hagg, T. and Robertson, H.A. (2004). Induction of neurogenesis in the adult rat subventricular zone and neostriatum following dopamine D receptor stimulation. *Eur. J. Neurosci.* 19(9): 2377–2387.
- Van Lookeren Campagne, M. and Gill, R. (1998). Tumor-suppressor p53 is expressed in proliferating and newly formed neurons of the embryonic and postnatal rat brain: Comparison with expression of the cell cycle regulators p21Waf1/Cip1, p27Kip1, p57Kip2, p16Ink4a, cyclin G1, and the proto-oncogene Bax. *J. Comp. Neurol.* 397(2): 181–198.
- Vanderluit, J.L., Ferguson, K.L., Nikolettou, V., Parker, M., Ruzhynsky, V., Alexson, T., McNamara, S.M., Park, D.S., Rudnicki, M. and Slack, R.S. (2004). p107 regulates neural precursor cells in the mammalian brain. *J. Cell Biol.* 166(6): 853–863.
- Vescovi, A.L., Reynolds, B.A., Fraser, D.D. and Weiss, S. (1993). bFGF regulates the proliferative fate of unipotent (neuronal) and bipotent (neuronal/astroglial) EGF-generated CNS progenitor cells. *Neuron* 11(5): 951–966.

- Viti, J., Feathers, A., Phillips, J. and Lillien, L. (2003a). Epidermal growth factor receptors control competence to interpret leukemia inhibitory factor as an astrocyte inducer in developing cortex. *J. Neurosci.* 23(8): 3385–3393.
- Viti, J., Gulacsi, A. and Lillien, L. (2003b). Wnt regulation of progenitor maturation in the cortex depends on Shh or fibroblast growth factor 2. *J. Neurosci.* 23(13): 5919–5927.
- Von Waechter, R. and Jaensch, B. (1972). Generation times of the matrix cells during embryonic brain development: An autoradiographic study in rats. *Brain Res.* 46: 235–250.
- Wagner, J.P., Black, I.B. and DiCicco-Bloom, E. (1999). Stimulation of neonatal and adult brain neurogenesis by subcutaneous injection of basic fibroblast growth factor. *J. Neurosci.* 19(14): 6006–6016.
- Wallace, V.A. (1999). Purkinje-cell-derived Sonic hedgehog regulates granule neuron precursor cell proliferation in the developing mouse cerebellum. *Curr. Biol.* 9(8): 445–448.
- Wanaka, A., Milbrandt, J. and Johnson, Jr., E.M. (1991). Expression of FGF receptor gene in rat development. *Development* 111(2): 455–468.
- Wechsler-Reya, R.J. and Scott, M.P. (1999). Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron* 22(1): 103114.
- Weinmaster, G. (2000). Notch signal transduction: A real rip and more. *Curr. Opin. Genet. Dev.* 10(4): 363–369.
- Wenger, R.H. (2002). Cellular adaptation to hypoxia: O₂-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O₂-regulated gene expression. *Faseb. J.* 16(10): 1151–1162.
- Whittemore, S.R., Morassutti, D.J., Walters, W.M., Liu, R.H. and Magnuson, D.S. (1999). Mitogen and substrate differentially affect the lineage restriction of adult rat subventricular zone neural precursor cell populations. *Exp. Cell Res.* 252(1): 75–95.
- Xian, C.J. and Zhou, X.F. (1999). Roles of transforming growth factor-alpha and related molecules in the nervous system. *Mol. Neurobiol.* 20(2-3): 157–183.
- Xu, X., Weinstein, M., Li, C., Naski, M., Cohen, R.I., Ornitz, D.M., Leder, P. and Deng, C. (1998). Fibroblast growth factor receptor 2 (FGFR2)-mediated reciprocal regulation loop between FGF8 and FGF10 is essential for limb induction. *Development* 125(4): 753–765.
- Yamada, S., Taketomi, T. and Yoshimura, A. (2004). Model analysis of difference between EGF pathway and FGF pathway. *Biochem. Biophys. Res. Commun.* 314(4): 1113–1120.
- Yamaguchi, T.P., Harpal, K., Henkemeyer, M. and Rossant, J. (1994). *fgfr-1* is required for embryonic growth and mesodermal patterning during mouse gastrulation. *Genes Dev.* 8(24): 3032–3044.
- Yamazaki, S., Iwamoto, R., Saeki, K., Asakura, M., Takashima, S., Yamazaki, A., Kimura, R., Mizushima, H., Moribe, H., Higashiyama, S., Endoh, M., Kaneda, Y., Takagi, S., Itami, S., Takeda, N., Yamada, G. and Mekada, E. (2003). Mice with defects in HB-EGF ectodomain shedding show severe developmental abnormalities. *J. Cell. Biol.* 163(3): 469–475.
- Yang, K. and Cepko, C.L. (1996). Flk-1, a receptor for vascular endothelial growth factor (VEGF), is expressed by retinal progenitor cells. *J. Neurosci.* 16(19): 6089–6099.

- Ying, Q.L., Nichols, J., Chambers, I. and Smith, A. (2003). BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 115(3): 281–292.
- Yoshida, M., Suda, Y., Matsuo, I., Miyamoto, N., Takeda, N., Kuratani, S. and Aizawa, S. (1997). *Emx1* and *Emx2* functions in development of dorsal telencephalon. *Development* 124(1): 101–111.
- Yoshikawa, K. (2000). Cell cycle regulators in neural stem cells and postmitotic neurons. *Neurosci. Res.* 37(1): 1–14.
- Yoshikawa, Y., Fujimori, T., McMahon, A.P. and Takada, S. (1997). Evidence that absence of *Wnt-3a* signaling promotes neuralization instead of paraxial mesoderm development in the mouse. *Dev. Biol.* 183(2): 234–242.
- Yoshimura, S., Takagi, Y., Harada, J., Teramoto, T., Thomas, S.S., Waeber, C., Bakowska, J.C., Breakefield, X.O. and Moskowitz, M.A. (2001). FGF-2 regulation of neurogenesis in adult hippocampus after brain injury. *Proc. Natl. Acad. Sci. U. S. A.* 98(10): 5874–5879.
- Yousoufian, H., Longmore, G., Neumann, D., Yoshimura, A. and Lodish, H.F. (1993). Structure, function, and activation of the erythropoietin receptor. *Blood* 81(9): 2223–2236.
- Yu, X., Shacka, J.J., Eells, J.B., Suarez-Quian, C., Przygodzki, R.M., Beleslin-Cokic, B., Lin, C.S., Nikodem, V.M., Hempstead, B., Flanders, K.C., Costantini, F. and Noguchi, C.T. (2002). Erythropoietin receptor signalling is required for normal brain development. *Development* 129(2): 505–516.
- Yun, K., Fischman, S., Johnson, J., Hrabe de Angelis, M., Weinmaster, G. and Rubenstein, J.L. (2002). Modulation of the notch signaling by *Mash1* and *Dlx1/2* regulates sequential specification and differentiation of progenitor cell types in the subcortical telencephalon. *Development* 129(21): 5029–5040.
- Zagon, I.S. and McLaughlin, P.J. (1987). Endogenous opioid systems regulate cell proliferation in the developing rat brain. *Brain Res.* 412(1): 68–72.
- Zechner, D., Fujita, Y., Hulsken, J., Muller, T., Walther, I., Taketo, M.M., Crenshaw, 3rd, E.B., Birchmeier, W. and Birchmeier, C. (2003). *beta*-Catenin signals regulate cell growth and the balance between progenitor cell expansion and differentiation in the nervous system. *Dev. Biol.* 258(2): 406–418.
- Zhang, H., Vutskits, L., Pepper, M.S. and Kiss, J.Z. (2003). VEGF is a chemoattractant for FGF-2-stimulated neural progenitors. *J. Cell Biol.* 163(6): 1375–1384.
- Zhang, R., Zhang, L., Zhang, Z., Wang, Y., Lu, M., Lapointe, M. and Chopp, M. (2001). A nitric oxide donor induces neurogenesis and reduces functional deficits after stroke in rats. *Ann. Neurol.* 50(5): 602–611.
- Zhu, G., Mehler, M.F., Zhao, J., Yu Yung, S. and Kessler, J.A. (1999). Sonic hedgehog and BMP2 exert opposing actions on proliferation and differentiation of embryonic neural progenitor cells. *Dev. Biol.* 215(1): 118–129.
- Zigova, T., Pencea, V., Wiegand, S.J. and Luskin, M.B. (1998). Intraventricular administration of BDNF increases the number of newly generated neurons in the adult olfactory bulb. *Mol. Cell Neurosci.* 11(4): 234–245.
- Zindy, F., Cunningham, J.J., Sherr, C.J., Jøgal, S., Smeyne, R.J. and Roussel, M.F. (1999). Postnatal neuronal proliferation in mice lacking *Ink4d* and *Kip1* inhibitors of cyclin-dependent kinases. *Proc. Natl. Acad. Sci. U. S. A.* 96(23): 13462–13467.
- Zindy, F., Quelle, D.E., Roussel, M.F. and Sherr, C.J. (1997). Expression of the p16INK4a tumor suppressor versus other INK4 family members during mouse development and aging. *Oncogene* 15(2): 203–211.

TABLE 1.

| Extracellular factor | Effect | Reference |
|------------------------|------------------------------|----------------------------------------------------------------|
| FGF2 | ↑ proliferation | (Wagner <i>et al.</i> , 1999) |
| | ↑ neurogenesis | (Wagner <i>et al.</i> , 1999) |
| EGF | ↑ proliferation | (Kuhn <i>et al.</i> , 1997) |
| | ↓ neurogenesis | (Doetsch <i>et al.</i> , 2002) |
| TGF α | ↑ proliferation | (Cooper and Isacson 2004) |
| | ↑ migration | |
| IGF-1 | ↑ proliferation | (Arsenijevic <i>et al.</i> , 2001) |
| | ↑ neurogenesis | (Arsenijevic and Weiss 1998) |
| | ↑ survival | (Gago <i>et al.</i> , 2003) |
| BDNF | ↑ proliferation (p75) | (Zigova <i>et al.</i> , 1998) |
| | ↑ neurogenesis (TrkB) | (Pencea <i>et al.</i> , 2001) |
| | ↑ survival | (Kirschenbaum and Goldman 1995) |
| EPO | ↑ neurogenesis | (Shingo <i>et al.</i> , 2001) |
| VEGF | ↑ proliferation | (Jin <i>et al.</i> , 2002b) |
| | ↑ migration | (Zhang <i>et al.</i> , 2003) |
| HB-EGF | ↑ neurogenesis | (Jin <i>et al.</i> , 2002a) |
| Ephrins | ↑ proliferation | (Conover <i>et al.</i> , 2000) |
| | ↓ migration | |
| TNF α | ↑ proliferation | (Wu <i>et al.</i> , 2000) |
| BMP | ↓ proliferation | (Coskun and Luskin, 2001) |
| | ↑ self-renewal | (Ying <i>et al.</i> , 2003) |
| | ↑ gliogenesis | (Gross <i>et al.</i> , 1996) |
| | ↑ neurogenesis | (Li <i>et al.</i> , 1998; Panchison <i>et al.</i> , 2001) |
| Noggin | ↑ neurogenesis | (Lim <i>et al.</i> , 2000) |
| CNTF/LIF | ↑ self-renewal | (Shimazaki <i>et al.</i> , 2001) |
| | ↑ gliogenesis | (Bonni <i>et al.</i> , 1997; Rajian <i>et al.</i> , 1998) |
| | ↑ neurogenesis | (Emsley and Hagg 2003) |
| Shh | ↑ proliferation | (Charytoniuk <i>et al.</i> , 2002; Palma <i>et al.</i> , 2005) |
| Wnt (β catenin) | +FGF2 proliferation | ↑ (Viti <i>et al.</i> , 2003; Israsena <i>et al.</i> , 2004) |
| | -FGF2 neurogenesis | ↑ (Israsena <i>et al.</i> , 2004; Otero <i>et al.</i> , 2004) |
| Notch | variable | (Chambers <i>et al.</i> , 2001) |
| Tenascin C | ↑ proliferation | (Garcion <i>et al.</i> , 2004) |
| Serotonin | ↑ proliferation | (Banasr <i>et al.</i> , 2004) |
| | ↑ neurogenesis | |
| Dopamine | ↑ proliferation | (Coronas <i>et al.</i> , 2004) |
| | (Baker <i>et al.</i> , 2004) | |
| Opioids | ↑ neurogenesis | (Van Kampen <i>et al.</i> , 2004) |
| | ↓ proliferation | (Stiene-Martin <i>et al.</i> , 2001) |
| Others | sAPP | ↑ proliferation (Ohsawa <i>et al.</i> , 1999) |
| | Abeta | ↓ proliferation (Haughey <i>et al.</i> , 2002) |
| | | ↓ migration |
| | | ↑ apoptosis |

**shh* enhanced the mitogenic effect of EGF