

# 1

## The Study of Biological Rhythms

*“Dost thou love life? Then do not squander time,  
for that’s the stuff life is made of.”  
—Benjamin Franklin (1706–1790)  
American Statesman & Philosopher*

### Introduction

How could you tell if a person is alive or dead? Chances are that your answer will depend upon the presence or absence of a rhythm, the beating of the heart. Biological rhythms are inherent to life itself and can be detected by all the senses. We can see them, hear them, feel them, smell them, and we may even taste evidence of them (Figure 1.1). Perhaps the sense of time itself could be considered a sixth sense. In many ways we can sense how long a time has elapsed since some occurrence was last noted, as well as the time of day, time of month, and time of year from cues all around us (Hering, 1940; Binkley, 1990). Life moves in synchrony to the beat of clocks and calendars, some outside the body and some within the very cells of all living things. Rhythms are among the common strands from which the web of life itself is spun.

By definition, a *rhythm* is a change that is repeated with a similar pattern. Humans, like all other organisms that inhabit this Earth, have a rhythmic order underlying life. Actually, change, not constancy, is the norm for life and the rhythmic timing of change makes predictability a reality. For example, throughout each 24-h day, leaves change their orientation (Figure 1.2), human body temperature rises and falls (Figure 1.3), fungi time their sporulation (Figure 1.4), and activity levels fluctuate (Figure 1.5). These rhythmic changes of life represent only a small segment of an enormous network of biological rhythms, passed on from one generation to the next.

All known variables of life, be they levels of potassium ions in a cell, stages of sleep, or the opening and closing of flowers, have either directly or indirectly been found to display rhythms. Furthermore, adaptations of organisms for survival relative to geophysical cycles, such as the solar day, seasons, and tides, attest to the evolution of the genetic aspects of certain types of rhythmic timing. The objective




FIGURE 1.1. Rhythms can be detected by the five senses: *sight* (migrations, plant movements, insect emergence); *sound* (roosters crowing or hens clucking, nocturnal activity of rodents, cricket chirping); *touch* (feeling sleepy, taking a pulse or feeling an insect bite at certain times of the day); *smell* (aromas from plants); and *taste* (starch vs. sugar in apple or corn; availability of Palolo worms near Pacific Islands).




FIGURE 1.2. Change in orientation of velvetleaf (*Abutilon theophrasti*) under a 24-h light-dark (LD) schedule. The relative position of a leaf at different times is shown under a schedule of 13 h of light (09:00–22:00 h) followed by 11 h of darkness (dark bars). Inset illustrates position of the blade in degrees from horizontal (=0°). Data obtained by video monitor in our lab.

FIGURE 1.3. The changes in human body temperature during usual living conditions. Dashed lines (—) indicate time when temperature was not recorded during sleep (shaded area/horizontal dark bars). Subject = healthy man (RBS, age 20 years) (Sothorn, unpublished).





FIGURE 1.4. Fungi (*Neurospora crassa*) growth and spore formation is circadian-rhythmic in constant conditions. Rhythmic spore (conidia) formation in *Neurospora crassa* can be observed when cultured on a nutritive medium in a glass tube. Inoculation was at left end of the tube, with the start of spore formation repeating at 24-h intervals (shown as bumps about every 24 h). Inset photo courtesy of Van D. Gooch, University of Minnesota-Morris.




FIGURE 1.5. Comparison of activity in animals and position of leaves in plants over 24 h. Diurnal animals, such as humans and birds, are active during the day, while nocturnal animals, such as hamsters and rodents, are at rest (*top panel*). At night, the reverse happens and while the diurnal animals are at rest, the nocturnal animals, such as the hamster and rat, are active (*bottom panel*). Also, the orientation of leaves changes from the horizontal during the day to the vertical at night.

of this chapter is to briefly introduce the topic of biological rhythms by emphasizing the commonality, presence, and importance of the subject in nearly every facet of life, as well as its status as an integrating discipline of biology.

## A Time for Everything

The rhythmic nature of life influences the very existence of organisms, commencing before conception and extending beyond death. Rhythms may be the most ubiquitous, yet overlooked, phenomena of life (Luce, 1970). They are such an integral part of life that the absence or perturbation of specific oscillations (e.g., brain waves and heart beats) in humans and other animals is used in the practice of medicine to distinguish between life and death, as well as between illness and good health (Figures 1.6 and 1.7). How organisms respond to certain chemicals, be they drugs or house dust, insecticides or herbicides, or any of a long list of agents is often time dependent (Table 1.1), with effects more pronounced at certain times than others. In some cases, what may be beneficial at one time may be noneffective or lethal at another. Even the perception of pain, be it disease-related (e.g., from headache, arthritis) or experimentally induced (e.g., by heat or cold or electrical stimuli), often shows a daily rhythm (Labrecque, 1992).




FIGURE 1.6. Diagram illustrating (A) the human heart and the specialized cardiac conduction system composed of the sinus node (SA), atrioventricular node (AV), right and left atrial (RA and LA) chambers, and the right and left ventricles (RV and LV). A small electrical current propagates from the sinus node through the atrial muscle to the AV node, which, after a short delay (usually 0.12 s), is transmitted to the ventricles. The electrical events cause both atrial and ventricular contraction. To record the electrical signals from the heart, electrodes are placed on the surface of the body at specific locations (e.g., right arm and left leg). The signals collected in this manner are amplified, and recorded on charts. This recording is called an electrocardiogram (ECG). (B) A record of an electrical cycle from a single lead (pair of electrodes). P = atrial electrical activity (depolarization), QRS complex defines the depolarization of ventricles; and T wave signifies repolarization of ventricles. (C) Three panels of ECGs (provided by Ann Dunnigan) illustrating their use in the diagnosis of normal as well as abnormal rhythms. *Panel 1*: Normal sinus rhythm rate (rate 75 beats per minute [bpm]); *Panel 2*: abnormally fast regular rhythm at rate of  $>175$  bpm; *Panel 3*: Abnormally slow rhythm where atria do not communicate with ventricles, called complete heart block. In *Panel 3*, sinus node rate ("P") is normal (75 bpm), while ventricular rate is abnormal (40 bpm).




FIGURE 1.7. Diagram illustrating dorsal views of the brain (A) and scalp showing the position of electrodes (B), and two electroencephalograms (EEG). *Left:* Normal brain EEG measuring electrical potential difference with respect to time shows posterior “alpha rhythm.” *Right:* Abnormal EEG in a young patient with epilepsy shows abnormal high amplitude discharges during a seizure. EEG panels provided by Mark Koukkari.

The concept of timing in relation to life and death is not of recent origin, as can be noted in the biblical book of Ecclesiastes 3:2–3:

*To everything there is a season and a time to every purpose under heaven: a time to be born and a time to die; a time to plant and a time to pluck up that which is planted; a time to kill and a time to heal; a time to break down and a time to build up ...*

The rhythmic nature of birth<sup>1</sup> (Figure 1.8), morbidity (presence of disease), and mortality<sup>2</sup> are widely documented in the scientific literature. While these are but a few of the many examples of daily changes, there are seasonal differences as well, which show an annual rhythm in these same events (Figure 1.9). Even the common things that we are aware of, such as the migration of birds (Gwinner, 1977, 2003), or the less common events, such as the germination of seeds in certain species, display an annual rhythm (Bünning & Mussle, 1951).

### Three Rhythm Domains

Much of the early work on biological rhythms focused on cycles in which the *period* was 24 h. These rhythms are referred to as *circadian rhythms* because they reoccur with a period of 24 h during usual light–dark conditions, and can

<sup>1</sup> Results from a classic study which summarized rhythms in human natality based upon the hourly incidence of 207,918 spontaneous labors and 2,082,453 natural births (Smolensky et al., 1972) showed that the onset of labor and birth under usual circumstances was most frequent between midnight and 06:00 h.

<sup>2</sup> Sudden cardiac death (Willich et al., 1987) and cerebral infarction (stroke) or hemorrhage (Marshall, 1977) are more prevalent between midnight and 06:00 h.

TABLE 1.1. Examples of time-dependent responses of organisms to various chemicals.

| Chemical                                | Organism   | Comments  | Reference                 |
|---|------------|---|---------------------------|
| Ethanol                                 | Human      | Blood levels greater at 07:00 h; poorer performance after 19:00 and 23:00 h                         | Reinberg et al., 1975     |
| Histamine                               | Human      | Greater reaction of lungs or skin near midnight   | Reinberg et al., 1969     |
| House dust                              | Human      | Bronchial hyperreactivity greater during nighttime than daytime                                     | Gervais et al., 1977      |
| Lidocaine (anesthetic)                  | Human      | Duration of anesthesia in teeth and skin longer in mid-afternoon                                    | Reinberg & Reinberg, 1977 |
| Theophylline (asthmatic drug)           | Human      | Sustained-release dose at 20:00 h better than 08:00 h in controlling nocturnal dip in lung function | Reinberg et al., 1987     |
| Corticosteroids                         | Human      | Once-daily dosing of inhaled steroid between 15:00 and 17:30 h optimal for asthma control           | Pincus et al., 1997       |
| Pentobarbital (anesthetic)              | Rat        | Duration of anesthesia longer in early dark (activity)  | Scheving et al., 1968     |
| Pentobarbital (anesthetic)              | Mouse      | Maximal mortality in early dark (activity)  | Bruguerolle et al., 1988  |
| Arabinosyl cytosine (anticancer drug)   | Mouse      | Increased tolerance in mid-light (resting)  | Haus et al., 1972         |
| Dichlorvos (insecticide)                | Cockroach  | Greatest toxicity in early dark   | Eesa et al., 1987         |
| Bentazon (herbicide)                    | Velvetleaf | Less injury near middle of the light span   | Koukkari & Johnson, 1979  |
| Glyphosate and glufosinate (herbicides) | Weeds      | Maximum control at midday   | Martinson et al., 2002    |

continue with a period close to 24.0 h when the organism is isolated from external cues. Thus, periods of about (*circa*) a day (*dies*) are present when organisms are isolated from environmental 24-h cycles, such as the alternating light and darkness of the solar day, and/or changes in temperature. This “free-running” period is an important characteristic of a circadian rhythm. In addition, the period is relatively consistent over a range of temperatures (temperature compensation). These and other characteristics are discussed more extensively in Chapter 2 on General Features of Rhythms.

While the circadian rhythms represent the dominant cycle that has been studied relative to the activities of humans and other organisms, cycles having periods shorter or longer than circadian, such as 90-min or seasonal cycles, are also important. Biological cycles that have periods less than 20 h are called *ultradian rhythms*, while cycles with periods longer than 28 h are called *infradian rhythms*. Depending upon the variable, infradian periods can be measured in weeks, months, years (circannual), and longer. Collectively, these three rhythmic domains comprise a network or web of rhythmic oscillations that in many ways can be likened to the various chemical pathways that perform different functions, but occur simultaneously within the same organelle or cell.

The range of periods for biological rhythms is broad, extending from cycles that are measured in milliseconds to cycles that are over 100 years in length. Some




FIGURE 1.8. Comparison of the circadian (*circa* = about, *dies* = day) waveform for the onset of labor and birth in humans, illustrating a natural sequence of events beginning with a peak in the onset of labor early in the night (dark bar) followed by a peak in natural births early in the morning. Hourly incidence of onset of >200,000 spontaneous labors and >2,000,000 natural births (redrawn from Smolensky et al., 1972).

examples illustrating the diversity in periods and types of variables are presented in Table 1.2 (ultradian and circadian domains) and Table 1.3 (infradian domain). Multiyear cycles, such as the emergence of the periodical cicada (*Magicicada* spp.) every 13 years in the south and midwestern USA or every 17 years in the northeastern USA, the 8- to 10-year population cycles of the ruffed grouse (*Bonasa umbellus*) in Minnesota, and the 15- to 120-year cycles in flowering of various species of bamboo, are among the spectacular rhythmic events found in nature that are little understood.




FIGURE 1.9. Monthly cardiac mortality shows circannual rhythm in the northern and southern hemispheres. A greater proportion of overall annual deaths from heart disease occurs in the winter for each hemisphere (redrawn from Smolensky et al., 1972).



TABLE 1.2. Examples of variables illustrating a range of rhythmic periods in the ultradian and circadian domains.

| Time    | Period     | Variable                                 | Organism                                      | Reference                  |
|---------|------------|--|---|----------------------------|
| Seconds | <1 s       | EEG activity (delta frequency)           | Human ( <i>Homo sapiens</i> )                 | Krippke, 1972              |
|         | <1 s       | ECG (depolarization of heart ventricles) | Human ( <i>Homo sapiens</i> )                 | see Figure 1.5             |
|         | 30–45 s    | NADH levels                              | Yeast ( <i>Saccharomyces carlsbergensis</i> ) | Chance et al., 1964        |
| Minutes | 1 min      | Courtship song (male)                    | Fruit fly ( <i>Drosophila melanogaster</i> )  | Kyriacou & Hall, 1980      |
|         | 2–4 min    | Leaflet movements                        | Telegraph plant ( <i>Desmodium gyrans</i> )   | Koukkari et al., 1985      |
|         | 4–15 min   | Reaction time                            | Human ( <i>Homo sapiens</i> )                 | Lovett-Douse et al., 1978  |
|         | 15 min     | Cortisol secretion                       | Horse (mares) ( <i>Equus caballus</i> )       | Drake & Evans, 1978        |
|         | 30 min     | Transpiration                            | Oat ( <i>Avena sativa</i> )                   | Johnsson, 1973             |
|         | 45–62 min  | Leaflet movements                        | Soybean ( <i>Glycine max</i> )                | Guillaume & Koukkari, 1987 |
|         | 76 min     | Cellular protein content                 | <i>Acanthamoeba castellanii</i>               | Lloyd et al., 1982         |
|         | 75–100 min | Pupillary motility                       | Human ( <i>Homo sapiens</i> )                 | Lavie, 1979                |
|         | 85–90 min  | Cortisol secretion                       | Monkey ( <i>Macaque mulatta</i> )             | Holaday et al., 1977       |
|         | 90 min     | Deep body and udder temperatures         | Holstein cow ( <i>Bos taurus</i> )            | Bitman et al., 1984        |
|         | 90–100 min | REM-NREM sleep                           | Human ( <i>Homo sapiens</i> )                 | Aserinsky & Kleitman, 1953 |
|         | 100 min    | Shoot movements (circumnutation)         | Pole bean ( <i>Phaseolus vulgaris</i> )       | Millet et al., 1984        |
| Hour    | 4 h        | Enzyme activity                          | Euglena ( <i>Euglena gracilis</i> )           | Balzer et al., 1989        |
|         | 12 h       | Amylase activity                         | Alfalfa ( <i>Medicago sativa</i> )            | Henson et al., 1986        |
| Day     | 24 h       | Body temperature                         | Human ( <i>Homo sapiens</i> )                 | Aschoff et al., 1972       |
|         | 24 h       | Sleep–wakefulness                        | Human ( <i>Homo sapiens</i> )                 | Kleitman, 1963             |
|         | 24 h       | Leaf movements                           | Albizzia ( <i>Albizzia julibrissin</i> )      | Koukkari, 1974             |
|         | 24 h       | Activity                                 | Flying squirrel ( <i>Galucomys volans</i> )   | DeCoursey, 1960            |

TABLE 1.3. Examples of variables illustrating a range of rhythmic periods in the infradian domain.

| Time  | Period             | Variable                   | Organism   | Reference                   |
|-------|--------------------|----------------------------|--|-----------------------------|
| Week  | 7 days             | Oviposition (egg laying)   | Spring Tail ( <i>Folsomia candida</i> )  | Chiba et al., 1973          |
|       | 7 days             | Organ transplant rejection | Human ( <i>Homo sapiens</i> )  | DeVecchi et al., 1981       |
|       | 7 days             | Imbibition of seeds        | Bean ( <i>Phaseolus vulgaris</i> )   | Spruyt et al., 1987         |
| Month | 27–34 days         | Menstrual cycle            | Woman ( <i>Homo sapiens</i> )  | Presser, 1974               |
|       | 6 months           | Ulcer perforation          | Human ( <i>Homo sapiens</i> )  | Svanes et al., 1998         |
| Year  | 1 year             | Seed germination           | <i>Digitalis lutea</i> , <i>Chrysanthemum corymbosum</i> , and <i>Gratiola officinalis</i> | Bünning & Müssle, 1951      |
|       |                    |                            | Wild potato ( <i>Solanum acaule</i> )  | Ludwig et al., 1982         |
|       |                    |                            | Pole bean ( <i>Phaseolus vulgaris</i> )  | Spruyt et al., 1988         |
|       | 1 year             | Antler replacement         | Sika deer ( <i>Cervus nippon</i> )   | Goss, 1969                  |
|       | 1 year             | Migration                  | Willow warbler (and others) ( <i>Phylloscopus trochilus</i> )                              | Gwinner, 1977               |
|       | 1 year             | Hibernation                | Golden-mantled ground squirrel ( <i>Citellus lateralis</i> )                               | Pengelley & Fisher, 1977    |
|       | 1 year             | Ovarian weight             | Yamuna River catfish ( <i>Heteropneustes fossilis</i> )                                    | Sundararaj et al., 1982     |
|       | 1 year and 7 years | Gonadal weight             | Purple sea urchin ( <i>Strongylocentrotus purpuratus</i> )                                 | Halberg et al., 1987        |
|       | 8 years            | Alpha-amylase activity     | Wheat ( <i>Triticum</i> spp.)  | Kettlewell et al., 1999     |
|       | 8–10 years         | Population                 | Ruffed Grouse ( <i>Bonasa umbellus</i> )   | Gullion, 1982, 1985         |
|       | 13 or 17 years     | Emergence                  | Periodical Cicada ( <i>Magicicada</i> spp.)  | Hoppensteadt & Keller, 1976 |
|       | 100–120 years      | Flowering                  | Chinese bamboo <sup>a</sup> ( <i>Phyllostachys bambusoides</i> )                           | Janzen, 1976                |

<sup>a</sup>While the range between successive synchronized reproductions by seed is from 3 to 120 years, most bamboo species have shorter, yet still very long times between seeding (15–60 years).

## Implications of Body Clocks

We have become a “clock driven” society, one that arranges time according to the demands of a commercial or industrial complex often fostered by profit and/or leisure, rather than arranging time so that our internal body clocks are in synchrony with the natural environment of this “clockwork Earth.” Accidents, catastrophes, and illnesses are inevitable when the time cycle of society does not heed the biological rules that underlie the rhythms of humans or other organisms.<sup>3</sup> For example, the reason that many traffic accidents occur during night and early morning hours is not only due to the difficulty in seeing in reduced light, but also to a decline in the alertness of the driver who is trying to overcome the physiological urge to sleep. Alertness is but one of the many performance variables that are under rhythmic control (Figure 1.10). It may be no coincidence that some of the major industrial catastrophes of our time were not related to weather, but were associated with erratic work–rest schedules and fatigue (see Essay 1.1).

### *Essay 1.1 (by RBS): Accidents and Catastrophes*

Four major industrial accidents that gained worldwide attention were each attributed to human error due to shift-work and its accompanying fatigue. The methyl isocyanate chemical accident in Bhopal, India, nuclear accidents at 3-Mile Island in the USA and in Chernobyl in the Ukraine, and the oil spill from the tanker Exxon Valdez in Alaska, all occurred at night (Reinberg & Smolensky, 1985; Moore-Ede, 1993). The nuclear accidents resulting in the release of radioactive gases occurred at 3-Mile Island at 04:00 h on March 28, 1979 and at 01:23 h in Chernobyl on April 26, 1986. The Exxon Valdez accident resulting in an environmentally catastrophic oil spill, occurred at 00:04 h on March 24, 1989, while the gas leak in Bhopal occurred at 00:56 h on December 3, 1984. All of these accidents occurred after midnight and were determined to be the result of operator error due to poor work–rest schedules, monotony, and fatigue. Interestingly, while hundreds of villagers and thousands of cattle in Bhopal died as they slept when the gas cloud swept over them, night-shift workers in the chemical plant, and nocturnally active rats observed scurrying around dead bodies all survived with little, if any, ill-effects from the toxic gas. Presumably, due to internal circadian time (stage) and not the external clock time, they were less susceptible to the toxic effects of the gas during the activity portion of their circadian sleep–wake cycle: rats are naturally active during the night, while the workers were in the active portion of their shifted circadian cycle.

Another less well-known, but major disaster occurred in western Canada on February 8, 1986 at 08:41 h, when a freight train going 94 km/h collided head-on with a passenger train going 78 km/h, killing 23 people and causing over \$30 million in damages (Smiley, 1990). Weather was not a factor, nor was there any hardware error; a red light was ignored by the engineer of the freight train and no brake application was made. The night before the accident, the freight engineer slept only 3.5 h, while the brakeman slept only 5 h before their train departed at 06:25 AM. Both had worked irregular schedules during

---

<sup>3</sup> This statement does not refer to the popular pseudoscience notion that good days and bad are based upon birth date (see discussion on Biorhythm Theory in Chapter 10).




FIGURE 1.10. Circadian patterns in some performance tests by a young man (RBS, 23 years of age) doing self-measurements every 2–4 h during waking-only for 2 weeks reveal that mental and physical performance were at their best in the middle of the daily activity span. Two-hour averages computed from a total of 90 measurements. Dark bar and shaded area indicate times of sleep when no data were obtained.

the previous month, with the engineer finishing a 6-h run at 23:30 h the night before. It was concluded that the number one factor underlying this catastrophe was human error brought on by the interaction between 24-h biologic rhythms and lengthy, irregular work–rest schedules, and consequent negative effects on alertness and performance of the freight crew.

The broad and interesting topic of the applications and implications of rhythms in society as concerns both humans and organisms in their environment will be discussed in Chapter 10 on Society.

## Chronobiology: An Integrating Discipline

The study of *biological rhythms*, known as *chronobiology*, is an integrating discipline that has been ranked parallel with the more classical disciplines of

TABLE 1.4. A comparison among the four disciplines of biology relative to duration and episodic designation (cf. Ehret, 1980).

| Discipline         | Period or duration          | Episodic designation |
|--------------------|-----------------------------|----------------------|
| Biological rhythms | Less than a second to years | Existence            |
| Development        | Seconds to years            | Life span            |
| Genetics           | Minutes to years            | Dynasty              |
| Evolution          | Hours to years              | Age                  |

*development*, *genetics*, and *evolution* (Ehret, 1980). All four of these disciplines span the structural levels of organization from molecules to ecosystems. A comparison among the four disciplines in regard to duration and episodic designation is presented in Table 1.4. The inclusion of chronobiology in formulating questions and hypotheses that lend themselves to experimentation now provides a broad unifying approach that extends from single-celled organisms to higher plants and animals. This can be seen by the continuing annual increase in the number of published scientific papers, which are accruing in the thousands per year since the late 1960s, and can be found in an online literature search<sup>4</sup> that uses circadian and other rhythm domain terms as key words (Figure 1.11).

Historically, the interdisciplinary nature of chronobiology was evident even in the composition of the small group of seven individuals who met in August 1937 in Sweden to create the first international organization focusing on the study of biological rhythms.<sup>5</sup> Of the three review papers presented at this conference that were of a more theoretical nature, one was based upon plant rhythms (by Anthonia Kleinhoonte from Holland), the second focused on animal rhythms (by Hans Kalmus from England), and the third dealt with human rhythms (by Arthur Jores from Germany) (Jores, 1938, 1975) (cf. Kalmus, 1974).<sup>6</sup>

<sup>4</sup> The online search engine PubMed (<http://www.ncbi.nlm.nih.gov/PubMed/>) was used for this literature search of articles indexed at the National Center for Biotechnology Information of the National Library of Medicine. There are hundreds of additional papers on rhythms in books, meeting proceedings, and journals, some of which can be found with other online search engines (e.g., Medline, BIOSIS, AGRICOLA).

<sup>5</sup> The next meeting of the International Society for the Study of Biological Rhythms was held with 12 participants in Holland just before World War II. The third meeting took place in Hamburg, Germany in 1949, with 50 participants. This organization changed its name to the International Society for Chronobiology at its first meeting in the USA in 1971 and included several hundred participants. There are now numerous societies studying rhythmic phenomena, including the American Association of Medical Chronobiology and Chronotherapeutics; the European Pineal Society; the European Society for Chronobiology; Groupe d'Etude des Rythmes Biologiques; Societa Italiana di Cronobiologia; the Japanese Society for Chronobiology, the Society for Light Treatment and Biological Rhythms; the Society for Research on Biological Rhythms; the Mediterranean Society for Chronobiology; among others.

<sup>6</sup> For historical notes see *Bulletin du Groupe d'Etude des Rythmes Biologiques* (1989) 21(1) (1er trimestre); and Cambrosio A, Keating P. (1983) *Social Studies of Science* 13: 323–353.




FIGURE 1.11. The number of articles on biological rhythms found by an online search engine (PubMed) using the key words circadian, diurnal, circannual, and ultradian, now number in the thousands over spans of 4 years. There are hundreds of additional papers on rhythms in books, meeting proceedings, and journals, some of which can be found with other online search engines (e.g., Medline, BIOSIS, AGRICOLA, etc.).

## Chapters in This Book

In this book, chapters have been prepared around main topics on biological rhythms. However, in some cases there is considerable overlap of material among chapters, which is unavoidable due to the ubiquity of the topic. Biological rhythms have a number of unique characteristics and terms that will be introduced in a separate chapter, and referred to throughout the book. Some of these characteristics, such as the stability of the period of circadian rhythms and how their phase<sup>7</sup> can be shifted, have been incorporated into models to better understand the mechanisms of timing (see Chapter 5 on Models) and how they contribute to the temporal organization of life. Clocks, calendars, and units of time that arose from cycles in the natural environment are described and compared with internal biological clocks in Chapter 3 on Time.

The rhythmic interactions between living things and the things that make them live are complex. Some are easily observed and/or measured, whereas others can be examined only indirectly by specific assays, images, or mathematical interpretations. How the rhythmic nature of life is best observed and monitored, as well as how data can be analyzed for rhythm characteristics, represent chapter topics (autorhythmometry, analyzing data) that help provide a basic understanding of biological rhythms. Because there is often a direct relationship between the internal rhythms of an organism and the external rhythms of the environment, the topic of photoperiodism, which focuses upon the response of

<sup>7</sup> Any point on a rhythmic pattern relative to a fixed reference point can be a phase. The time of a peak or trough relative to local midnight are two commonly used phases. In many laboratory studies, the zero point coincides with the start of the light span (e.g., circadian time), rather than midnight.

organisms (be they plant or animal) to given durations and times of light and darkness over the seasons, is included as a chapter. Chapters that deal with tidal and lunar rhythms, sexuality and reproduction, natural resources and agriculture, society, and clinical and veterinary medicine, describe from a practical perspective how knowledge of biological rhythms applies to, or affects, nearly all aspects of life.

## Take-Home Message

Biological rhythms are common phenomena of life and are found in all major groups of organisms, but are often overlooked. The length of time required to repeat a rhythmic cycle is called the period, a characteristic that has been used to categorize rhythms into three major groups: circadian (20–28 h), ultradian (<20 h), and infradian (>28 h). Circadian rhythms are usually synchronized by cyclic changes in light and darkness and/or temperature. When isolated from such environmental cues, the rhythm continues (free-runs), but usually with a period slightly longer or shorter than precisely 24.0 h. Some typical examples of biological variables for rhythms include (a) *circadian*: body temperature in humans and leaf movements of plants; (b) *ultradian*: brain waves of humans and twining of movements of bean shoots; and (c) *infradian*: the menstrual cycle of human females and the annual germination of certain seeds.

## References

- Aschoff J, Giedke H, Pöppel E, Wever R. (1972) The influence of sleep-interruption and of sleep-deprivation on circadian rhythms in human performance. In: *Aspects of Human efficiency. Diurnal Rhythm and Loss of Sleep*. Colquhoun WP, ed. London: English University Press, pp. 133–150.
- Aserinsky E, Kleitman N. (1953) Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. *Science* 118(3062): 273–274.
- Balzer I, Neuhaus-Steinmetz U, Quentin E, van Wüllen M, Hardeland R. (1989) Concomitance of circadian and circa-4-hour ultradian rhythms in *Euglena gracilis*. *J Interdiscipl Cycle Res* 20: 15–24.
- Binkley S. (1990) *The Clockwork Sparrow*. Englewood Cliffs, NJ: Prentice-Hall, 262 pp.
- Bitman J, Lefcourt A, Wood DL, Stroud B. (1984) Circadian and ultradian temperature rhythms of lactating dairy cows. *J Dairy Sci* 67(5): 1014–1023.
- Bruguerolle B, Prat M, Douylliez C, Dorfman P. (1988) Are there circadian and circannual variations in acute toxicity of phenobarbital in mice? *Fundam Clin Pharmacol* 2(4): 301–304.
- Bünning E, Müsle L. (1951) Der Verlauf der endogen Jahresrhythmik in Samen unter dem Einfluss verschiedenartiger Aussenfaktoren. *Z Naturforsch* 6b: 108–112.
- Chance B, Estabrook RW, Ghosh A. (1964) Damped sinusoidal oscillations of cytoplasmic reduced pyridine nucleotide in yeast cells. *Proc Natl Acad Sci (USA)* 51: 1244–1251.
- Chiba Y, Cutkomp LK, Halberg F. (1973) Circaseptan (7-day) oviposition rhythm and growth of Spring Tail, *Folsomia Candida* (Collembola: Isotomidae). *J Interdiscipl Cycle Res* 4: 59–66.

- DeCoursey PJ. (1960) Phase control of activity in a rodent. In: *Biological Clocks. Cold Spring Harbor Symposia on Quantitative Biology*, Vol 25. New York: Long Island Biol Assoc, pp. 49–55.
- DeVecchi A, Halberg F, Sothorn RB, Cantaluppi A, Ponticelli C. (1981) Circaseptan rhythmic aspects of rejection in treated patients with kidney transplant. In: *Chronopharmacology and Chronotherapeutics*. Walker CA, Winget CM, Soliman KFA, eds. Tallahassee: Florida A&M University Foundation, pp. 339–353.
- Drake DJ, Evans JW. (1978) Cortisol secretion pattern during prolonged ACTH infusion in dexamethasone treated mares. *J Interdiscipl Cycle Res* 9: 89–96.
- Eesa N, Cutkomp LK, Cornélissen G, Halberg F. (1987) Circadian change in Dichloros lethality (LD50) in the cockroach in LD 14:10 and continuous red light. In: *Advances in Chronobiology, Part A*. Pauly JE, Scheving LE, eds. New York: Alan R. Liss, pp. 265–279.
- Ehret CF. (1980) On circadian cybernetics, and the innate and genetic nature of circadian rhythms. In: *Chronobiology: Principles and Applications to Shifts in Schedules*. Scheving LE, Halberg F, eds. Alphen aan den Rijn: Sijthoff & Noordhoff, pp. 109–125.
- Gervais P, Reinberg A, Gervais C, Smolensky MH, De France O. (1977) Twenty-four-hour rhythm in the bronchial hyperreactivity to house dust in asthmatics. *J Allergy Clin Immunol* 59(3): 207–213.
- Goss RJ. (1969) Photoperiodic control of antler cycles in deer: I. Phase shift and frequency changes. *J Exp Zool* 170: 311–324.
- Guillaume FM, Koukkari WL. (1987) Two types of high frequency oscillations in *Glycine max* (L.) Merr. In: *Advances in Chronobiology, Part A*. Pauly JE, Scheving LE, eds. New York: Alan R. Liss, pp. 47–58.
- Gullion GW. (1982) Forest wildlife interactions. In: *Introduction to Forest Science*. Young RA, ed. New York: Wiley, pp. 379–407.
- Gullion GW. (1985) Ruffed grouse research at the University of Minnesota Cloquet Forestry Center. *Minn Dept Nat Res Wildl Res Unit 1985 Report*, pp. 40–49.
- Gwinner E. (1977) Circannual rhythms in bird migration. *Annu Rev Ecol Syst* 8: 381–405.
- Gwinner E. (2003) Circannual rhythms in birds. *Curr Opin Neurobiol* 13(6): 770–778.
- Halberg Fcn, Halberg F, Sothorn RB, Pearse JS, Pearse VB, Shankaraiah K, Giese AC. (1987) Consistent synchronization and circaseptennian (about 7-yearly) modulation of circannual gonadal index rhythm of two marine invertebrates. In: *Advances in Chronobiology—Part A*. Pauly JE, Scheving LE, eds. New York: Alan R. Liss, pp. 225–238.
- Haus E, Halberg F, Scheving LE, Pauly JE, Cardoso S, Kühl JFW, Sothorn RB, Shiotsuka RN, Hwang DS. (1972) Increased tolerance of leukemic mice to arabinosyl cytosine with schedule adjusted to circadian system. *Science* 177(43): 80–82.
- Henson CA, Duke SH, Koukkari WL. (1986) Rhythmic oscillations in starch concentration and activities of amylolytic enzymes and invertase in *Medicago sativa* nodules. *Plant Cell Physiol* 27: 233–242.
- Hering DW. (1940) The time concept and time sense among cultured and uncultured peoples. In: *Time and Its Mysteries, Series II*. New York: New York University Press, pp. 3–39.
- Holaday JW, Martinez HM, Natelson BH. (1977) Synchronized ultradian cortisol rhythms in monkeys: persistence during corticotropin infusion. *Science* 198(4312): 56–58.
- Hoppensteadt FC, Keller JB. (1976) Synchronization of periodical cicada emergences. *Science* 194(4262): 335–337.
- Janzen DH. (1976) Why bamboos wait so long to flower. *Annu Rev Ecol Syst* 7: 347–391.
- Johnsson A. (1973) Oscillatory transpiration and water uptake of Avena plants: I. Preliminary observations. *Physiol Plant* 28: 40–50.



- Jores A. (1938) First Conf. Ronneby, Sweden, August 13–14, 1937. *Dtsch Med Wochenschr* 64(21/28): 737–989.
- Jores A. (1975) The origins of chronobiology: an historical outline. *Chronobiologia* 2(2): 155–159.
- Kalmus H. (1974) The foundation meeting of the international society for biological rhythms. *Chronobiologia* 1: 118–124.
- Kettlewell PS, Sothorn RB, Koukkari WL. (1999) U.K. wheat quality and economic value are dependent on the North Atlantic Oscillation. *J Cereal Science* 29: 205–209.
- Kleitman N. (1963) *Sleep and Wakefulness*, 2nd edn, Chicago, IL: University of Chicago Press.
- Koukkari WL. (1974) Rhythmic movements of *Albizia julibrissin* pinnules. In: *Chronobiology*. Scheving LE, Halberg F, Pauly JE, eds. Tokyo: Igaku Shoin, pp. 676–678.
- Koukkari WL, Johnson MA. (1979) Oscillations of leaves of *Abutilon theophrasti* (velvetleaf) and their sensitivity to bentazon in relation to low and high humidity. *Physiol Plant* 47: 158–162.
- Koukkari WL, Duke SH, Bonzon MV. (1985) Circadian rhythms and their relationships to ultradian and high frequency oscillations. In: *Les Mécanismes de l'Irritabilité et du Fonctionnement des Rythmes chez les Végétaux, 1977–1983*. Grippin H, Wagner E, eds. Genève: Université de Genève, pp. 106–126.
- Kripke DF. (1972) An ultradian biological rhythm associated with perceptual deprivation and REM sleep. *Psychosom Med* 34(3): 221–234.
- Kyriacou CP, Hall JC. (1980) Circadian rhythm mutations in *Drosophila melanogaster* affect short-term fluctuations in the male's courtship song. *Proc Natl Acad Sci USA* 77(11): 6729–6733.
- Labrecque G. (1992) Inflammatory reaction and disease states. In: *Biological Rhythms in Clinical and Laboratory Medicine*. Touitou Y, Haus E, eds. Berlin: Springer-Verlag, pp. 483–492.
- Lavie P. (1979) Ultradian rhythms in alertness—a pupillometric study. *Biol Psychol* 9(1): 49–62.
- Lloyd D, Edwards SW, Fry JC. (1982) Temperature-compensated oscillations in respiration and cellular protein content in synchronous cultures of *Acanthamoeba castellanii*. *Proc Natl Acad Sci (USA)* 79(12): 3785–3788.
- Lovett-Douse JW, Payne WD, Podnieks I. (1978) An ultradian rhythm of reaction time measurements in man. *Neuropsychobiology* 4(2): 93–98.
- Luce G. (1970) *Biological Rhythms in Psychiatry and Medicine*. Washington, DC: Natl Inst Mental Health, US Dept. Health, Education and Welfare, 183 pp.
- Ludwig H, Hinze E, Junges W. (1982) Endogene Rhythmen des Keimverhaltens der Samen von Kartoffeln, insbesondere von *Solanum acaule*. *Seed Sci Technol* 10: 77–86.
- Marshall J. (1977) Diurnal variation in the occurrence of strokes. *Stroke* 8(2): 230–231.
- Martinson KB, Sothorn RB, Koukkari WL, Durgan BR, Gunsolus JL. (2002) Circadian response of annual weeds to Glyphosate and Glufosinate. *Chronobiol Intl* 19(2): 405–422. [Erratum: *Chronobiol Intl* 2002; 19(4): 805–806]
- Millet B, Melin D, Bonnet B, Ibrahim CA, Mercier J. (1984) Rhythmic circumnutation movement of the shoots in *Phaseolus vulgaris* L. *Chronobiol Intl* 1(1): 11–19.
- Moore-Ede M. (1993) *The Twenty-Four Hour Society: Understanding Human Limits in a World That Never Stops*. Reading, MA: Addison-Wesley, 230 pp.
- Pengelley ET, Fisher KC. (1963) The effect of temperature and photoperiod on the yearly hibernating behavior of captive golden-mantled ground squirrels (*Citellus lateralis testicorum*). *Can J Zool* 41: 1103–1120.

- Pincus DJ, Humeston TR, Martin RJ. (1997) Further studies on the chronotherapy of asthma with inhaled steroids: the effect of dosage timing on drug efficacy. *J Allergy Clin Immunol* 100(6 Pt 1): 771–774.
- Presser HB. (1974) Temporal data relating to the human menstrual cycle. In: *Biorhythms and Human Reproduction*. Ferin M, Halberg F, Richert RM, Vande Wiele R, eds. New York: Wiley, pp. 145–160.
- Reinberg A, Zagula-Mally Z, Ghata J, Halberg F. (1969) Circadian reactivity rhythm of human skin to house dust, penicillin, and histamine. *J Allergy* 44(5): 292–306.
- Reinberg A, Clench J, Aymard N, Galliot M, Bourdon R, Gervais P, Abulker C, Dupont J. (1975) [Circadian variations of the effects of ethanol and of blood ethanol values in the healthy adult man. Chronopharmacological study] [French]. *J Physiol (Paris)* 70(4): 435–456.
- Reinberg A, Reinberg MA. (1977) Circadian changes in the duration of local anesthetic agents. *Naunyn-Schmiedeberg's Arch Pharmacol* 297: 149–152.
- Reinberg A, Smolensky MH. (1985) Chronobiologic considerations of the Bhopal methyl isocyanate disaster. *Chronobiol Intl* 2(1): 61–62.
- Reinberg A, Pauchet F, Ruff F, Gervais A, Smolensky MH, Levi F, Gervais P, Chaouat D, Abella ML, Zidani R. (1987) Comparison of once-daily evening versus morning sustained-release theophylline dosing for nocturnal asthma. *Chronobiol Intl* 4(3): 409–419.
- Scheving LE, Vedral D, Pauly JA. (1968) Circadian susceptibility rhythm in rats to pentobarbital sodium. *Anat Rec* 160(4): 741–750.
- Smiley AM. (1990) The Hinton train disaster. *Accid Anal Prev* 22(5): 443–455.
- Smolensky M, Halberg F, Sargent F. (1972) Chronobiology of the life sequence. In: *Advances in Climatic Physiology*. Ito S, Ogata K, Yohimura H, eds. Tokyo: Igaku Shoin, pp. 281–318.
- Spruyt E, Verbelen J-P, De Greef JA. (1987) Expression of circaseptan and circannual rhythmicity in the imbibition of dry stored bean seeds. *Plant Physiol* 84: 707–710.
- Spruyt E, Verbelen J-P, De Greef JA. (1988) Ultradian and circannual rhythmicity in germination of *Phaseolus* seeds. *J Plant Physiol* 132: 234–238.
- Sundararaj BI, Vasal S, Halberg F. (1982) Circannual rhythmic ovarian recrudescence in the catfish, *Heteropneustes fossilis* (Bloch). In: *Toward Chronopharmacology*. Takahashi R, Halberg F, Walker C, eds. New York: Pergamon, pp. 319–337.
- Svanes C, Sothorn RB, Sørbye H. (1998) Rhythmic patterns in incidence of peptic ulcer perforation over 5.5 decades in Norway. *Chronobiol Intl* 15(3): 241–264.
- Willich SN, Levy D, Rocco MB, Tofler GH, Stone PH, Muller JE. (1987) Circadian variation in the incidence of sudden cardiac death in the Framingham Heart Study population. *Amer J Cardiol* 60: 801–806.