

# Propagation Through Trapped Sets and Semiclassical Resolvent Estimates

Kiril Datchev and András Vasy

Let  $P = -h^2 \Delta + V(x)$ ,  $V \in C_0^\infty(\mathbb{R}^n)$ . We are interested in semiclassical resolvent estimates of the form

$$\|\chi(P - E - i0)^{-1} \chi\|_{L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)} \leq \frac{a(h)}{h}, \quad h \in (0, h_0], \quad (1)$$

for  $E > 0$ ,  $\chi \in C^\infty(\mathbb{R}^n)$  with  $|\chi(x)| \leq \langle x \rangle^{-s}$ ,  $s > 1/2$ . We ask: how is the function  $a(h)$  for which (1) holds affected by the relationship between the support of  $\chi$  and the trapped set at energy  $E$ , defined by

$$K_E = \{\alpha \in T^*\mathbb{R}^n : \exists C > 0, \forall t > 0, |\exp(tH_p)\alpha| \leq C\}?$$

Here  $p = |\xi|^2 + V(x)$  and  $H_p = 2\xi \cdot \nabla_x - \nabla V \cdot \nabla_\xi$ .

We have (1) with  $\chi(x) = \langle x \rangle^{-s}$  and  $a(h) = C$  for all  $E$  in a neighborhood of  $E_0 > 0$  if and only if  $K_{E_0} = \emptyset$  ([6, 7]). For general  $V$  and  $\chi$ , the optimal bound is  $a(h) = \exp(C/h)$ , but Burq [1] and Cardoso-Vodev [2] prove that for any given  $V$ , if  $\chi$  vanishes on a sufficiently large compact set, for any  $E > 0$  there exists  $C$  such that (1) holds with  $a(h) = C$ . In our main theorem we improve the condition on  $\chi$  and obtain a shorter proof at the expense of an a priori assumption.

**Theorem 1 ([3]).** Fix  $E > 0$ . Suppose that (1) holds for  $\chi(x) = \langle x \rangle^{-s}$  with  $s > 1/2$  and with  $a(h) = h^{-N}$  for some  $N \in \mathbb{N}$ . Then if we take instead  $\chi$  such that  $K_E \cap T^* \text{supp } \chi = \emptyset$ , we have (1) with  $a(h) = C$ .

---

K. Datchev (✉)

Mathematics Department, MIT, Cambridge, MA, 02139-4307, USA

e-mail: [datchev@math.mit.edu](mailto:datchev@math.mit.edu)

A. Vasy

Mathematics Department, Stanford University, Stanford, CA, 94305-2125, USA

e-mail: [andras@math.stanford.edu](mailto:andras@math.stanford.edu)

In fact our result holds for more general operators, and the cutoff  $\chi$  can be replaced by a cutoff in phase space whose microsupport is disjoint from  $K_E$ . In certain situations it is even possible to take a cutoff whose support overlaps  $K_E$ : see [3] for more details and references.

The a priori assumption that (1) holds for  $\chi(x) = \langle x \rangle^{-s}$  with  $a(h) = h^{-N}$  is not present in [1, 2] and is not always satisfied, but there are many examples of hyperbolic trapping where it holds: see e.g. [5, 8].

To indicate the comparative simplicity of our method, we prove a special case of the Theorem, under the additional assumption that  $\text{supp } V \subset \{|x| < R_0\}$  and  $\text{supp } \chi \subset \{R_0 < |x| < R_0 + 1\}$ . In other words, suppose  $(P - \lambda)u = f$ , with  $\text{Re } \lambda = E$ , and  $\text{supp } f \subset \{R_0 < |x| < R_0 + 1\}$ ,  $\|f\| \leq 1$ . We must prove that  $\|\chi u\| \leq Ch^{-1}$ , uniformly as  $\text{Im } \lambda \rightarrow 0^+$ . Here and below all norms are  $L^2$  norms.

Let  $S$  denote functions in  $C^\infty(T^*\mathbb{R}^n)$  which are bounded together with all derivatives, and for  $a \in S$  define

$$\text{Op}(a)u(x) = (2\pi h)^{-n} \int \exp(i(x - y) \cdot \xi/h) a(x, \xi) u(y) dy d\xi.$$

Because  $P - \lambda$  has a semiclassical elliptic inverse away from  $p^{-1}(E)$  (see for example [4, Chap. 4]), we have  $\|\text{Op}(a)u\| \leq C$  whenever  $\text{supp } a \cap p^{-1}(E) = \emptyset$ . Consequently it is enough to show that  $\|\text{Op}(a)u\| \leq Ch^{-1}$  for some  $a \in S$  with  $a$  nowhere vanishing on  $T^* \text{supp } \chi \cap p^{-1}(E)$ . We will prove this inductively: we will show that if there is  $a_1$  with this nowhere vanishing property such that  $\|\text{Op}(a_1)u\| \leq Ch^k$ , then there is  $a_2$  with the same nowhere vanishing property such that  $\|\text{Op}(a_2)u\| \leq Ch^{k+1/2}$ , provided  $k \leq -3/2$ . The base case follows from the a priori assumption that  $\|u\| \leq h^{-N-1}$ , so it suffices to prove the inductive step.

Take  $\varphi = \varphi(|x|) \geq 0$  a smooth function such that  $\varphi = 1$  when  $|x| \leq R_0$ ,  $\varphi = 0$  when  $|x| \geq R_0 + 1$ ,  $\varphi' = -\psi^2$  with  $\psi$  smooth. We require further that  $T^* \text{supp } \psi$  be contained in the set where  $a_1$  is nonvanishing, and in the end we will take  $a_2 = \psi$ . We will now use a positive commutator argument with  $\varphi$  as the commutant:

$$i\langle [P, \varphi]u, u \rangle = i\langle u, \varphi f \rangle - i\langle \varphi f, u \rangle - 2\text{Im } \lambda \|u\|^2 \geq -C\|\psi u\|\|f\|, \quad (2)$$

where we used first  $(P - \lambda)u = f$  and then  $\text{Im } \lambda \geq 0$  and  $\text{supp } f \subset \{\psi \neq 0\}$ . The semiclassical principal symbol of  $i[P, \varphi]$  is

$$hH_p\varphi = 2h\rho\varphi' = -2h\rho\psi^2,$$

where  $\rho$  is the dual variable to  $|x|$  in  $T^*\mathbb{R}^n$ .

We now define an open cover and partition of unity of  $T^* \text{supp } \chi$  according to the regions where this commutator does and does not have a favorable sign (the favorable sign is  $H_p\varphi < 0$ , because of the direction of the inequality in (2)). Take  $c > 0$  small enough that for  $\rho < 2c$ ,  $|x| > R_0$ ,  $t < 0$  we have  $x + 2\rho t \notin \text{supp } V$ . Let  $K$  be a neighborhood of  $p^{-1}(E) \cap T^* \text{supp } \chi$  with compact closure in  $T^*\{R_0 < |x| < R_0 + 1\}$ , and let  $O$  be a neighborhood of  $K$  with compact closure

in  $T^*\{R_0 < |x| < R_0 + 1\}$ , and let

$$U_+ = \{\alpha \in O : \rho > c\}, \quad U_- = \{\alpha \in O : \rho < 2c\} \cup (T^*\mathbb{R}^n \setminus K).$$

Take  $\phi_\pm \in C_0^\infty(O)$  with  $\phi_+^2 + \phi_-^2 = 1$  on  $T^*\text{supp } \chi$  and with  $\text{supp } \phi_\pm \subset U_\pm$ .

Then

$$H_p \varphi = -b^2 - 2\rho\psi^2\phi_-^2, \quad \text{where } b = \sqrt{2\rho}\psi\phi_+,$$

and if  $B = \text{Op}(b)$  and  $\Phi_- = \text{Op}(\phi_-)$

$$i[P, \varphi] = -hB^*B + h\Phi_-R_1\Phi_- + h^2R_2 + O(h^\infty),$$

where  $R_{1,2} = \text{Op}(r_{1,2})$  for  $r_{1,2} \in S$  with  $\text{supp } r_{1,2} \subset \text{supp } \psi$ . Combining with (2), and using  $L^2$  boundedness of  $R_1$ , we obtain

$$h\|Bu\|^2 \leq Ch\|\Phi_-u\|^2 + h^2\langle R_2u, u \rangle + C\|\psi u\|\|f\| + O(h^\infty).$$

Since  $\langle R_2u, u \rangle \leq Ch^{2k}$  by inductive hypothesis, we have

$$\begin{aligned} \|Bu\|^2 &\leq C(\|\Phi_-u\|^2 + h^{2k+1} + h^{-1}\|\psi u\|\|f\|) \\ &\leq C(\|\Phi_-u\|^2 + h^{2k+1} + \delta^{-1}h^{-2} + \delta\|\psi u\|^2), \end{aligned}$$

where we used  $\|f\| \leq 1$ , and where  $\delta > 0$  will be specified presently. Since at least one of  $B$  and  $\Phi_-$  is elliptic at each point in the interior of  $T^*\text{supp } \psi$ , we have

$$\|\psi u\|^2 \leq C(\|\Phi_-u\|^2 + \|Bu\|^2), \quad (3)$$

from which we conclude that, if  $\delta$  is sufficiently small,

$$\|Bu\|^2 \leq C_\delta(\|\Phi_-u\|^2 + h^{-2} + h^{2k+1}). \quad (4)$$

Because  $c$  was chosen small enough that all backward bicharacteristics through  $\text{supp } \phi_-$  stay in  $T^*\{|x| > R_0\}$ , where  $P = -h^2\Delta$ , we have

$$\|\Phi_-u\| \leq Ch^{-1},$$

by standard nontrapping estimates (see, for example, [3, Sect. 6]). This, combined with (3) and (4), gives

$$\|\psi u\|^2 \leq C_\delta(h^{-2} + h^{2k+1}),$$

after which taking  $a_2 = \psi$  completes the proof of the inductive step.

**Acknowledgements** The first author is partially supported by a National Science Foundation postdoctoral fellowship, and the second author is partially supported by the NSF under grant DMS-0801226, and a Chambers Fellowship from Stanford University. The authors are grateful for the hospitality of the Mathematical Sciences Research Institute, where part of this research was carried out.

## References

1. Nicolas Burq. Lower bounds for shape resonances widths of long range Schrödinger operators. *Amer. J. Math.*, 124:4, 677–735, 2002.
2. Fernando Cardoso and Georgi Vodev. Uniform estimates of the resolvent of the Laplace-Beltrami operator on infinite volume Riemannian manifolds II. *Ann. Henri Poincaré*, 3:4, 673–691, 2002.
3. Kiril Datchev and András Vasy. Propagation through trapped sets and semiclassical resolvent estimates. To appear in *Ann. Inst. Fourier*. Preprint available at arXiv:1010.2190.
4. Lawrence C. Evans and Maciej Zworski. Lecture notes on semiclassical analysis. Available online at <http://math.berkeley.edu/~zworski/semiclassical.pdf>.
5. Stéphane Nonnenmacher and Maciej Zworski. Quantum decay rates in chaotic scattering. *Acta Math.* 203:2, 149–233, 2009.
6. Didier Robert and Hideo Tamura. Semiclassical estimates for resolvents and asymptotics for total scattering cross-sections. *Ann. Inst. H. Poincaré Phys. Théor.* 46:4, 415–442, 1987.
7. Xue Ping Wang. Semiclassical resolvent estimates for  $N$ -body Schrödinger operators. *J. Funct. Anal.* 97:2, 466–483, 1991.
8. Jared Wunsch and Maciej Zworski. Resolvent estimates for normally hyperbolic trapped sets. To appear in *Ann. Inst. Henri Poincaré (A)*. Preprint available at arXiv:1003.4640.



<http://www.springer.com/978-3-0348-0465-3>

Microlocal Methods in Mathematical Physics and Global  
Analysis

Grieser, D.; Teufel, S.; Vasy, A. (Eds.)

2013, IX, 148 p. 2 illus. in color., Softcover

ISBN: 978-3-0348-0465-3

A product of Birkhäuser Basel